

2 Admitted patient characteristics

The focus (for admitted patients) of this report is the data reported to the NHMD for the NMDS—Admitted Patient Mental Health Care, i.e. admitted patient separations that included a component of specialised psychiatric care. This first section reports these data within the context of 'mental health-related separations', i.e. all separations either with a mental health principal diagnosis or that included a component of specialised psychiatric care.

A mental health principal diagnosis in this publication is defined as any separation whose principal diagnosis falls within the chapter on mental disorders in the ICD-10-AM classification (F00 to F99), and some other diagnoses (Appendix 1). When comparing the 1998–99 figures with the previous year, note that codes for post-partum depression which were included in *Disorders of Pregnancy, childbirth and the puerperium* in ICD-9-CM have been moved to the mental disorders chapter in ICD-10-AM and are therefore included in the definition of a mental health principal diagnosis for 1998–99 data (see Appendix 2).

Data are also presented on separations with specialised psychiatric care in the Major Diagnostic Category (MDC of the AR-DRG 4.0/4.1 classification) related to mental diseases and disorders. This section also includes some comparisons between the mental health-related and the specialised psychiatric care separations.

Overview

Mental health-related separations

Table 2.1 summarises the general and specialised psychiatric care in 1998–99 in terms of same day separations and overnight separations and patient days (see Glossary for definition of overnight separations). In 1998–99, mental health-related separations accounted for 4.4% of total hospital separations (251,428 separations) and 14.2% of all patient days (3,177,971 days). In 1997–98, mental health-related separations (which were defined to exclude post-partum depression) accounted for 4.4% of total hospital separations (245,583 separations) and 14.6% of all patient days (3,303,205 days) (AIHW: Moore et al. 2000).

During 1998–99, 37.9% of mental health-related separations (95,222 separations) were same day separations compared with 36.6% (89,866) for the 1997–98 collection period. Overnight separations in 1998–99 accounted for 97% of the mental health-related patient days (3,082,749 days). The proportion of overnight separation patient days for 1997–98 was also 97% (3,059,756).

Figure 2.1 illustrates the relationship between separations with specialised psychiatric care and separations with a mental health principal diagnosis within the context of all hospital separations for 1998–99. Of the 251,428 mental health-related separations reported for 1998–99, there were 168,579 separations or 67.1% with at least some specialised psychiatric care. In comparison, of the 245,583 mental health-related separations reported for 1997–98, there were 133,887 separations or 54.5% with at least some specialised psychiatric care.

Of the 235,487 separations with a mental health principal diagnosis reported for 1997–98, there were 130,222 separations or 55.3% with at least some specialised psychiatric care. In comparison, of the 239,930 separations with a mental health principal diagnosis reported for

1998–99, there were 157,081 separations or 65.5% with at least some specialised psychiatric care. There were 82,849 separations with a mental health principal diagnosis that did not record a component of specialised psychiatric care for 1998–99. The equivalent figure for 1997–98 was 111,696 separations.

Of the separations with specialised psychiatric care, 157,081 or 93.2% were reported as having a mental health principal diagnosis in 1998–99. In 1997–98, a majority of specialised psychiatric care separations (123,791 separations or 92.5%) were also reported as having a mental health principal diagnosis. Thus, there was little change over the year in the proportion of separations that have a mental health principal diagnosis within the specialised separations group.




Figure 2.1: Separations with a mental health principal diagnosis and separations with specialised psychiatric care, Australia, 1998–99

Figure 2.2 outlines the relationship between patient days relating to separations with specialised psychiatric care and those with a mental health principal diagnosis. It shows that patient days of separations with some specialised psychiatric care accounted for nearly 80% of all patient days of separations with a mental health principal diagnosis (2,392,599 days with some specialised psychiatric care of a total 2,999,416 patient days).

In 1998–99 the total number of mental health-related patient days was 3,177,971 (see Table 2.1). Of these, 97% or 3,082,749 days were in overnight separations. Approximately 73% of total mental health-related patient days were from overnight separations with both a mental health principal diagnosis and specialised psychiatric care; 5.5% were attributable to overnight separations with specialised psychiatric care but no mental health principal diagnosis. Overnight separations with a mental health principal diagnosis but no specialised psychiatric care accounted for 18.4% of all mental health-related patient days.




Figure 2.2: Patient-days with a mental health principal diagnosis and patient-days in specialised psychiatric care, Australia, 1998–99

Table 2.2 provides the number of mental health-related separations and patient days for each State and Territory.

Table 2.3 divides mental health-related separations into those with and without specialised psychiatric care and outlines the principal diagnosis of each separation. It provides the summary information illustrated in Figures 2.1 and 2.2. A more detailed discussion of principal diagnoses for separations with specialised psychiatric care can be found in Section 2.2.

Major Diagnostic Category—mental diseases and disorders

This section examines the relationship between separations with specialised psychiatric care and those in the MDC relating to mental diseases and disorders (MDC 19).

AR-DRGs categorise acute admitted patient episodes into groups with similar clinical conditions and resource requirements (see Appendix 2 for further information on AR-DRG Version 4). These categories were designed to provide clinically meaningful way of relating the number and type of patients treated in a hospital (that is, its casemix), to the resources required by the hospital. AR-DRG categories can be grouped into MDCs and used to compare aspects of hospitalisation such as length of stay. Note that the data presented by AR-DRGs and MDCs include non-acute admitted patient episode data.




Figure 2.3 shows that of all separations with specialised psychiatric care, 90.2% were in MDC 19. Table 2.4 outlines all separations which fall into MDC 19 (Mental diseases and disorders) and MDC 20 (Alcohol & drug-related disorders) or have specialised psychiatric care. Of the 179,297 separations in MDC 19, 138,884 or 77.5% received some specialised psychiatric care. MDC 19 separations accounted for a total of 2,149,506 patient days, of which 76.5% were spent in specialised psychiatric care. In comparison, in the MDC related to alcohol and other drugs (MDC 20), 49.3% of all separations received some specialised psychiatric care and 56% of patient days were psychiatric care days.

One third of total separations with specialised psychiatric care were classified in AR-DRG U60Z; *Mental health treatment, same day, without electroconvulsive therapy*. This group accounted for 78.6% of all same day separations in the NMDS—Admitted Patient Mental Health Care data collection.

Of the 97,065 overnight separations, 23,623 (24.3%) were classified as belonging to the two AR-DRGs associated with principal diagnoses of *Schizophrenia disorders* (12,441 with involuntary legal status and 11,182 without involuntary legal status). The next largest group of overnight separations was *Major affective disorders age < 70 without catastrophic or severe complications and comorbidities* (U63B) with 21,367 separations.

Major affective disorders age < 70 without catastrophic or severe complications and comorbidities (U63B) was the AR-DRG with the highest number of separations of those in the admitted patient mental health collection (24,651 separations, 21,367 with specialised psychiatric care), but *Schizophrenia disorders with involuntary mental health legal status* (U61A) contributed the highest number of patient days and psychiatric care days (619,334 and 486,296 days respectively).

Separations with specialised psychiatric care

In 1998–99, separations with specialised psychiatric care accounted for 2.9% of total hospital separations (168,579 separations) and 11.5% of all patient days (2,571,154 days). In 1997–98, separations with specialised psychiatric care accounted for 2.4% of total hospital separations (133,887 separations) and 10.7% of all patient days (2,421,460 days) (AIHW: Moore et al. 2000).

In 1998–99, 42.2% of separations with specialised psychiatric care (71,514 separations) were same day separations compared with 41.6% (55,807 separations) for 1997–98. Overnight separations in 1998–99 accounted for 97.2% of the patient days (2,499,640 days).

Table 2.5 summarises the separations, patient days and psychiatric care days in 1998–99 of all separations with specialised psychiatric care by hospital type and jurisdiction. Public acute hospitals accounted for 62.0% of the total number of overnight specialised separations among the three hospital types. Over 70% of specialised separations from public acute hospitals were overnight separations. Private hospitals dominated the same day specialised separations with 62.0% of the total number of same day separations. Approximately 69% of specialised psychiatric care separations in private hospitals were same day. Public psychiatric hospitals had the greatest number of reported patient days and psychiatric care days for specialised psychiatric care separations, with 49.0% of the total number of patient days and 44.1% of the total number of psychiatric care days.

Principal diagnoses for admitted patients

This section outlines the principal diagnoses of separations with specialised psychiatric care collected from State and Territory governments under the NMDS—Admitted Patient Mental Health Care. The data presented in this section detail hospital separations in which specialised psychiatric care was reported.

Principal diagnosis is presented in this report in ICD-10-AM, but note that Queensland, Western Australia, South Australia and Tasmania provided data for 1998–99 using the ICD-9-CM classification. Data provided in ICD-9-CM were mapped to ICD-10-AM by AIHW (see Appendix 5).

This section presents data on the principal diagnoses using groupings within the ICD-10-AM chapter on mental disorders, but it also presents data in subsets of these groupings in a way similar to the reporting of mental disorders for National Health Priority Areas (AIHW 2000). This finer level of detail will affect the relative ranking of disorders.

Table 2.6 shows the distribution of separations with specialised psychiatric care principal diagnosis. In 1998–99, *Depressive disorders* (F32–F33) accounted for the highest number of same day separations (24,041) and overnight separations (19,181), and the highest per capita rate of hospitalisation at 2.3 hospital separations per 1,000 population. Almost 34% of all same day separations and 20% of all overnight separations had principal diagnoses in the *Depressive disorders* (F32–F33) category. The second largest grouping of principal diagnoses for same-day separations was *Neurotic, stress-related and somatoform disorders* (F40–48), reported for 13,436 same day separations (18.8%). *Schizophrenia* (F20) had the second highest number of overnight separations (18,311) and was reported for 18.9% of all overnight separations.

Schizophrenia (F20) accounted for the greatest number of patient days and psychiatric care days with 871,557 or 33.9% of patient days and 692,839 or 31.9% of total psychiatric care days. The number of psychiatric care days for *Schizophrenia* (F20) was approximately double

the number of days attributable to *Depressive disorders* (F32–F33) (369,308 patient days and 328,939 psychiatric care days). This was reflected in the figures for average length of stay (overnight) where *Schizophrenia* (F20) had 47.4 patient days per specialised separation and 37.6 psychiatric care days per separation, whereas *Depressive disorders* (F32–F33) had 18 and 16 days respectively. The principal diagnosis categories of *Mental retardation* (F70–79) and *Disorders of psychological development* (F80–89) had relatively small numbers of separations but much higher rates of patient days per separation, i.e. length of stay, than *Schizophrenia* (F20) and *Depressive disorders* (F32–F33) (see Table 2.6).

Table 2.7 shows the number of separations with specialised psychiatric care by principal diagnosis in ICD-10-AM chapter groupings. Almost 93% of the separations with specialised psychiatric care had a principal diagnosis from the ICD-10-AM mental and behavioural disorders chapter, and they accounted for 107.4 psychiatric care days per 1,000 population. Specialised psychiatric care separations with a principal diagnosis outside the ICD-10-AM mental health chapter accounted for 7.2% of specialised separations and 6.9% of psychiatric care days.

Hospital type

The distribution of specialised separations by principal diagnosis for each hospital type is presented in Table 2.8. Separations in the principal diagnosis group of *Depressive disorders* (F32–F33) accounted for approximately 38% of all private hospital separations and 41% of private hospital psychiatric care days. Separations with principal diagnoses of *Neurotic, stress-related and somatoform disorders* (F40–F48) accounted for 18.8% of all private hospital separations and 13.9% of private hospital psychiatric care days. Same day separations with principal diagnoses of *Depressive disorders* (F32–F33) made up 38.6% of private same day separations, and same day separations with principal diagnoses of *Neurotic, stress-related and somatoform disorders* (F40–F48) made up 20.5% of private same day separations. For overnight separations, the corresponding figures were 37.6% and 15.0% respectively.

Approximately 20% of all public acute hospital separations had a principal diagnosis in the group *Depressive disorders* (F32–F33), which accounted for 18.1% of reported public acute hospital psychiatric care days. Almost 18% of public acute hospital separations were separations with a principal diagnosis of *Schizophrenia* (F20), which accounted for 31.5% of reported public acute psychiatric care days. Separations with a principal diagnosis of *Schizophrenia* (F20) accounted for 22.5% of public acute hospital specialised overnight separations, and *Depressive disorders* (F32–F33) had the second highest proportion with 17.0%.

Over 22% of all public psychiatric hospital separations and 41.0% of all psychiatric care days in public psychiatric hospitals were attributed to a principal diagnosis of *Schizophrenia* (F20). *Schizophrenia* (F20) also accounted for the highest proportion of overnight separations and patient days in public psychiatric hospitals (22.4% and 41.0% respectively). The principal diagnosis group *Mental and behavioural disorders due to other psychoactive substances use* (F11–F19) accounted for the highest proportion of same day separations in public psychiatric hospitals (21.4%).

States and Territories

Tables 2.9 to 2.12 outline the number of separations, patient days and psychiatric care days in each principal diagnosis group for each jurisdiction for the 1998–99 financial year. State and Territory data relate to the State and Territory of the hospital, not the State or Territory

of the patient's usual residence. These figures can be influenced by a number of factors such as:

- the availability of admitted patient specialised psychiatric care in the jurisdiction;
- the spread of the population in rural, remote and metropolitan areas;
- differing admission practices; and
- the availability of community-based psychiatric facilities.

Some differences may also be attributed to the mapping of diagnosis codes from ICD-9-CM to ICD-10-AM for Queensland, Western Australia, South Australia and Tasmania (see Appendix 5).

Rates were indirectly age-standardised using the total separations, patient days or psychiatric care days for 1998–99 and the estimated resident population as at December 1998 to calculate the expected number of separations for each jurisdiction (see Appendix 4). Indirect age-standardisation has been used throughout this publication where indicated because this method is less sensitive to small cell sizes than the direct standardisation method.

Table 2.9 presents same day separations with specialised psychiatric care for all States and Territories. Over half the same day separations for Australia as a whole were for principal diagnoses in either *Depressive disorders* (F32–F33) (33.6% of total same day separations) or *Neurotic, stress-related and somatoform disorders* (F40–48) (18.8% of total same-day separations). *Depressive disorders* (F32–F33) accounted for the highest number of same day separations for each jurisdiction except South Australia and the two Territories. The principal diagnosis group *Neurotic, stress-related and somatoform disorders* (F40–48) was the second largest category of same day separations after *Depressive disorders* (F32–F33) in New South Wales, Queensland and Western Australia. Conversely, *Neurotic, stress-related and somatoform disorders* (F40–48) was the largest category of same day separations in South Australia (26.0%), with *Schizophrenia* (F20) being the second largest category (20.8%) followed closely by *Depressive disorders* (F32–F33) (19.9%). In Victoria the second largest category of same day separations was *Eating disorders* (F50), and in Tasmania it was *Bipolar affective disorders* (F31). Western Australia had the highest age-standardised rate of same day separations (7.4 separations per 1,000 population), nearly twice the rates of New South Wales (3.75) and Victoria (4.1). This compares with an age-standardised rate for Australia overall of 3.8 separations per 1,000 population.

Overnight separations with specialised psychiatric care are presented in Table 2.10. For Australia as a whole, *Depressive disorders* (F32–F33) and *Schizophrenia* (F20) were the principal diagnoses contributing the greatest number of overnight separations (19.7% and 18.9% respectively). *Depressive disorders* (F32–F33) contributed the highest number of overnight separations in Victoria, Queensland, Western Australia and South Australia. In Western Australia the number of overnight separations reported for *Depressive disorders* (F32–F33) was almost matched by *Neurotic, stress-related and somatoform disorders* (F40–48) and *Schizophrenia* (F20). A principal diagnosis of *Schizophrenia* (F20) contributed the highest number of overnight separations in New South Wales, Tasmania, the Australian Capital Territory and Northern Territory. Queensland had the highest age-standardised rate of overnight separations per 1,000 population at 6.7, compared with an Australian rate of 5.2.

Separations with a principal diagnosis of *Schizophrenia* (F20) accounted for the highest number of patient days as well as the highest number of specialised psychiatric care days for all jurisdictions (Tables 2.11 and 2.12). Separations in the principal diagnosis group *Depressive disorders* (F32–F33) accounted for the next highest number of patient days and psychiatric care days for New South Wales, Victoria, Queensland, South Australia and

Western Australia (patient days only), and *Bipolar affective disorders* (F31) had the second highest number of patient days and psychiatric care days for Tasmania and the Australian Capital Territory. In the Northern Territory *Other schizophrenic, schizotypal, delusional disorders* (F21–29) had the second highest number of patient days and psychiatric care days, and in the Australian Capital Territory numbers in this category were the third highest following closely behind *Bipolar affective disorders* (F31).

The indirectly age-standardised rate for patient days per 1,000 population and psychiatric care days per 1,000 population were comparatively high in Queensland (223.9 patient days and 221.7 psychiatric care days per 1,000 population), almost twice the national rate (132.7 patient days and 106.8 psychiatric care days per 1,000 population). The Australian Capital Territory and the Northern Territory had the lowest rates at 31.5 and 46.4 patient days per 1,000 population respectively and 31.5 and 42.7 psychiatric care days per 1,000 population. This is possibly because these Territories have no public psychiatric hospitals, and this could affect the type of diagnoses treated in admitted patient settings in these Territories, as well as the amount of specialised psychiatric care provided.

Demographic profile of admitted patients with specialised psychiatric care

This section presents the socio-demographic data collected from State and Territory governments under the NMDS—Admitted Patient Mental Health Care. The section includes data for 6 of the 9 socio-demographic data elements in the NMDS; *Age, Sex, Marital status, Area of usual residence, Aboriginal and Torres Strait Islander status* and *Country of birth*. The data that were assessed to be of poor quality or inconsistently applied among jurisdictions were for *Employment status, Pension status* and *Type of usual accommodation* and these were excluded. Chapter 5 provides details on the data developments anticipated for these excluded data elements.

The data presented in this section detail separations in which specialised psychiatric care was reported and reflects the level of utilisation of hospital services by specific population groups. It does not necessarily reflect the prevalence of mental disorders in the broader community or the utilisation of community-based services by specific population groups.

Age and sex

Figure 2.4 presents the rates of separations per 1,000 population for each age and sex category. Males aged 25–34 years and females aged 35–44 years had the highest number of separations with specialised psychiatric care per 1,000 population. Females had higher separation rates than males for all age groups from age 18 to 64 years.

Tables 2.13a and b and 2.14a and b detail the distribution of separations with specialised psychiatric care by age group, sex and principal diagnosis. Male patients accounted for 43.4% of same day separations with specialised psychiatric care. *Depressive disorders* (F32–F33) was the largest principal diagnosis category for male patients, constituting 27.2% of male same day separations (8,435 separations). The second largest category was *Neurotic, stress-related and somatoform disorders* (F40–F48), which accounted for 23.9% of male same day separations (7,425). Relative to male patients, female patients had a higher proportion of same day separations with principal diagnoses of *Depressive disorders* (F32–F33) (38.6% or 15,606 separations). *Neurotic, stress-related and somatoform disorders* (F40–F48) accounted for 6,011 female separations (14.9%). The principal diagnosis of *Eating disorders* (F50), which was

reported for 111 same day separations for males, was the third highest number of reported same day separations for females (11.6% or 4,697 separations).

Male patients accounted for 50.2% of overnight separations (48,754) with specialised psychiatric care during 1998–99. *Schizophrenia* (F20) constituted 25.6% of male overnight separations, *Depressive disorders* (F32–F33) accounted for 14.2% and *Neurotic, stress-related and somatoform disorders*, 11.4%. For female patients, *Depressive disorders* (F32–F33) contributed the highest number of overnight separations (12,239 or 25.3%), followed by *Neurotic, stress-related and somatoform disorders* (F40–F48) (6,273 or 13%) and *Schizophrenia* (F20) (5,848 or 12.1%). A greater proportion of overnight separations in the principal diagnosis group of *Depressive disorders* (F32–F33) were reported for female patients, with 25.3% of overnight female separations in this category compared with 14.2% for male patients.

The number of male overnight separations with principal diagnoses of *Mental and behavioural disorders due to psychoactive substance use* (F10–F19) was more than double that for females. This difference was largest in the 18–34 years age groups for psychoactive substances and in the 25–54 years age groups for alcohol.

The proportion of male overnight separations with a principal diagnosis of *Schizophrenia* (F20) disorders (25.5%) was more than double the proportion and numbers for female patients (12.1%). For males, the number of these separations peaked in the 25–34 years age group with elevated levels also reported for the 18–24 years and 35–44 years age groups. For females the pattern was slightly different with the number of these separations also peaking in the 25–34 year age group but with elevated levels reported for the 35–44 years and 45–54 years age groups.

There were greater proportions of separations for principal diagnoses of *Neurotic, stress-related and somatoform disorders* (F40–F48) for females (13.0%) than for males (11.4%), with greater proportions reported for females in all age groups except the 45–64 years age group. The largest numbers of female and male overnight separations in the *Neurotic, stress-related and somatoform disorders* (F40–F48) category were in the 25–34 years age groups.

Overnight separations with principal diagnoses of *Depressive disorders* (F32–F33) accounted for 13.2% of separations of females aged 17 years and under. This proportion increased over age groups to 39.8% of separations of females in the 65 years and over category.

Approximately 8% of male overnight separations aged 17 years and under had a principal diagnosis in the *Depressive disorders* (F32–F33) category, and this proportion increased to 29.8% for those aged 65 years and over.

Overnight separations with a principal diagnosis of *Eating disorders* (F50) predominantly involved female patients. However, *Eating disorders* (F50) did not dominate the female overnight separations (2.1%) to the same extent as the female same day separations (11.6%). The number of separations with a principal diagnosis of *Eating disorders* (F50) was greatest in the younger age groups and decreased with age. The highest number of overnight separations in this category was reported in the 18–24 years age group.

Tables 2.15a and b and 2.16a and b show that the principal diagnosis of *Schizophrenia* (F20) was reported for the highest number of overnight patient days and psychiatric care days for both male and female patients. Principal diagnoses in the *Depressive disorders* (F32–F33) category had the second highest number of overnight patient days and psychiatric care days for both male and female patients.

Figure 2.5 shows the number of psychiatric care days per overnight separation for males and females for selected disorders. Although males had the highest proportion of overnight separations for the principal diagnosis of *Schizophrenia* (F20), the number of psychiatric care days per overnight separation was higher for females than for males. Female patients

received more than twice the number of psychiatric care days per overnight separation for *Manic episode* (F30) than male patients.

Marital status

Table 2.17 outlines separations with specialised psychiatric care by marital status, sex and jurisdiction. Marital status data for 1998–99 were assessed as unreliable for Victoria during the NMDS—Admitted Patient Mental Health Care validation process. Marital status data were unavailable for private hospitals in New South Wales and South Australia. Figure 2.6 presents the proportion of separations for male and female patients by marital status category after removing Victorian and private hospital data. With unreliable data removed, almost half (48.0%) of separations with specialised psychiatric care for males were for patients who had never been married, compared with just under one-third of female patients (31.5%). A further 21.3% of specialised psychiatric care separations were for male patients who were married or living in a de facto relationship, compared with 25.9% of female patients.

Area of usual residence

Table 2.18 demonstrates that, Australia-wide, patients from areas classified as remote or rural had fewer specialised separations per 1,000 population than patients from areas classified as metropolitan. In 1998–99 there were 10 separations of patients from metropolitan areas per 1,000 population compared with 1.7 separations per 1,000 population from rural areas and 2.6 separations per 1,000 population for patients from remote areas. In Western Australia and Northern Territory, the separation rates for patients from remote areas (13.1 and 3.5 respectively) were higher than separation rates for patients from metropolitan areas (9.4 and 2.1 respectively). In Tasmania the separation rate for rural areas (8.4 separations per 1,000 population) was higher than the metropolitan rate of 7.2 separations per 1,000 population. Queensland had the highest rate of specialised separations for patients from metropolitan areas (11.6 separations per 1,000 population), Tasmania for patients from rural areas (8.4 separations per 1,000 population), and Western Australia for remote areas (13.1 separations per 1,000 population).

Figures 2.7 and 2.8 show the number of psychiatric care days per overnight separation in each Rural, Remote and Metropolitan Area (RRMA) category of the patient's area of usual residence for public acute/private hospitals and public psychiatric hospitals respectively. Whereas private hospitals had higher numbers of psychiatric care days for metropolitan and rural areas, the length of psychiatric care in private hospitals for patients from remote areas was shorter than the length of psychiatric care for similar patients in public acute hospitals. In the private sector the highest overall average number of psychiatric care days per separation was for metropolitan areas other than capital cities.

Public psychiatric hospitals had higher numbers of psychiatric care days per overnight separation than either public acute or private separations. In public psychiatric hospitals the highest number of psychiatric care days per separation was for patients usually residing in rural areas, with an average of approximately 112 days for large rural centres and 96.7 days for other rural areas. Patients received an average of about 40 psychiatric care days per separation if they usually resided in a remote centre and an average of about 26 if they usually resided in another remote area.

Aboriginal and Torres Strait Islander status

Table 2.19 details the number of separations, patient days and psychiatric care days for Aboriginal and Torres Strait Islander peoples in 1998–99. The quality of Aboriginal and Torres Strait Islander status data varies, and so these figures should be used with caution. Further detail about the quality of these data can be found in *Australian Hospital Statistics 1998–99*.

During 1998–99, 383 same day and 2,678 overnight specialised separations were reported for Aboriginal and Torres Strait Islander peoples nationwide. Principal diagnoses of *Neurotic, stress-related and somatoform disorders* (F40–48) accounted for 42.6% of same day separations, and *Schizophrenia, schizotypal and delusional disorders* (F20–F29) accounted for the highest number of overnight separations (33.7%), patient days (60.7%) and specialised psychiatric care days (61.4%). Principal diagnoses of *Schizophrenia, schizotypal and delusional disorders* (F20–F29) accounted for 2.7 separations per 1,000 Aboriginal and Torres Strait Islander population (age-standardised). *Mental and behavioural disorders due to psychoactive substances use* (F10–F19) accounted for 17.7% of total separations at an age-standardised rate of 1.6 specialised separations per 1,000 Aboriginal and Torres Strait Islander population.

The specialised separation rate per 1,000 population for Aboriginal and Torres Strait Islander peoples (9.0 per 1,000 population) and the rate for all patients (8.9 per 1,000 population) were similar. The rate for Aboriginal and Torres Strait Islander peoples was notably higher than the total Australian specialised separation rate for principal diagnoses of *Schizophrenia, schizotypal and delusional disorders* (F20–F29) and *Mental and behavioural disorders due to psychoactive substances use* (F10–F19) (Figure 2.9). The rate for Aboriginal and Torres Strait Islander patients was much lower than the Australian specialised separation rate for *Mood (affective) disorders* (F30–F39).

There were, on average, 23 psychiatric care days per overnight separation for identified Aboriginal and Torres Strait Islander patients. There were, on average, 20.7 psychiatric care days per overnight separation reported for all patients. Figure 2.10 compares the rates of psychiatric care days per overnight separation for all patients and patients identified as Aboriginal and Torres Strait Islanders for selected principal diagnosis groups.

Table 2.20 presents the distribution of Aboriginal and Torres Strait Islander patients separations, patient days and psychiatric care days by area of usual residence, categorised using RRMA categories. Over half (56%) of the specialised separations reported for Aboriginal and Torres Strait Islander patients were for patients usually residing in metropolitan areas, compared with 83% for all patients.

Country of birth

Tables 2.21 and 2.22 detail the number of specialised separations, patient days and psychiatric care days for admitted patients according to their country of birth, grouped into Australia, Other English-speaking country and Non-English-speaking country (see Glossary).

Nationally, 6,141 same day and 7,556 overnight separations were reported for patients born in English-speaking countries other than Australia, and 11,583 same day and 17,378 overnight separations were reported for patients born in non-English-speaking countries. Principal diagnoses of *Mood (affective) disorders* (F30–F39) accounted for the highest number of specialised separations for all birth country groups. *Neurotic, stress-related and somatoform disorders* (F40–48) accounted for the second highest number of same day separations for all three groups and *Schizophrenia, schizotypal and delusional disorders* (F20–F29) accounted for

the second highest number of overnight separations and the highest number of patient days for all birth-country groups.

The number of patient days, specialised psychiatric care days and the crude rate of psychiatric care days per 1,000 population for the three country of birth categories are given in Table 2.22. Figure 2.11 illustrates the number of psychiatric care days per overnight separation for selected disorder categories by country of birth. *Schizophrenia* (F20) and *Manic episode* (F30) had the highest average number of specialised psychiatric care days per separation, but in both of these categories patients born in other English-speaking countries received fewer days in specialised psychiatric care on average. *Eating disorders* (F50) also had high levels of psychiatric care days per overnight separation for each country of birth category.

Table 2.1: Specialised and general care of all mental health-related separations^(a), Australia, 1998–99

	Number	Per cent of all hospital Separations	Per cent of all mental health-related
Same day separations			
All same day separations ^(b)	2,747,617	47.91	
With specialised psychiatric care			
with mental health principal diagnosis	66,235	1.15	26.34
without mental health principal diagnosis	5,279	0.09	2.10
Other mental health principal diagnosis without specialised care	23,708	0.41	9.43
<i>Total mental health-related same day separations</i>	<i>95,222</i>	<i>1.66</i>	<i>37.87</i>
Overnight separations			
All overnight separations ^(b)	2,987,432	52.09	
With specialised psychiatric care			
with mental health principal diagnosis	90,846	1.58	36.13
without mental health principal diagnosis	6,219	0.11	2.47
Other mental health principal diagnosis without specialised care	59,141	1.03	23.52
<i>Total mental health-related overnight separations</i>	<i>156,206</i>	<i>2.72</i>	<i>62.13</i>
Total mental health-related separations	251,428	4.38	100.00
Patient days			
All patient days			
All patient days due to same day separations ^(b)	2,747,617	12.31	
Patient days for same day separations			
With specialised psychiatric care			
with mental health principal diagnosis	66,235	0.30	2.08
without mental health principal diagnosis	5,279	0.02	0.17
Other mental health principal diagnosis without specialised care	23,708	0.11	0.75
<i>Total mental health-related same day patient days</i>	<i>95,222</i>	<i>0.43</i>	<i>3.00</i>
Patient days for overnight separations			
All patient days due to overnight separations ^(b)	19,571,424	87.69	
With specialised psychiatric care ^(c)			
with mental health principal diagnosis	2,326,364	10.42	73.20
without mental health principal diagnosis	173,276	0.78	5.45
Other mental health principal diagnosis without specialised care	583,109	2.61	18.35
<i>Total overnight patient days with a mental health principal</i>	<i>3,082,749</i>	<i>13.81</i>	<i>97.00</i>
Total mental health-related patient days	3,177,971	14.24	100.00

(a) Includes all separations with a mental health principal diagnosis as defined in Appendix 1 and all separations for which specialised psychiatric care was reported.

(b) From *Australian Hospital Statistics 1998–99*.

(c) Count of patient days for separations with some specialised psychiatric care.

Table 2.2: Specialised and general care of all mental health-related separations^(a), States and Territories, 1998–99

	NSW	Vic	QLD	WA	SA	Tas	ACT	NT	Total
Separations									
Same day separations									
With specialised psychiatric care									
with mental health principal diagnosis	22,054	18,450	10,749	12,036	2,137	745	30	34	66,235
without mental health principal diagnosis	1,848	660	1,174	1,509	12	66	6	4	5,279
Other mental health principal diagnosis without specialised care	7,981	6,282	4,165	1,422	1,358	2,325	110	65	23,708
<i>Total mental health-related same day separations</i>	31,883	25,392	16,088	14,967	3,507	3,136	146	103	95,222
Overnight separations									
With specialised psychiatric care									
with mental health principal diagnosis	27,640	21,702	18,895	9,484	9,740	1,954	739	692	90,846
without mental health principal diagnosis	1,786	1,370	1,485	818	261	415	52	32	6,219
Other mental health principal diagnosis without specialised care	18,739	13,384	11,759	6,827	5,556	1,817	855	204	59,141
<i>Total mental health-related overnight separations</i>	48,165	36,456	32,139	17,129	15,557	4,186	1,646	928	156,281
Total mental health-related separations	80,048	61,848	48,227	32,096	19,064	7,322	1,792	1,031	251,503
Patient days for same day separations									
With specialised psychiatric care									
with mental health principal diagnosis	22,054	18,450	10,749	12,036	2,137	745	30	34	66,235
without mental health principal diagnosis	1,848	660	1,174	1,509	12	66	6	4	5,279
Other mental health principal diagnosis without specialised care	7,981	6,282	4,165	1,422	1,358	2,325	110	65	23,708
<i>Total mental health-related same day patient days</i>	31,883	25,392	16,088	14,967	3,507	3,136	146	103	95,222
Patient days for overnight separations									
With specialised psychiatric care ^(b)									
with mental health principal diagnosis	766,444	354,566	718,014	237,667	189,868	43,003	9,091	7,711	2,326,364
without mental health principal diagnosis	63,894	25,763	51,434	8,228	4,005	19,250	418	284	173,276
Other mental health principal diagnosis without specialised care	200,193	117,173	135,510	50,424	43,417	21,553	13,526	1,313	583,109
<i>Total overnight patient days with a mental health principal diagnosis</i>	1,030,531	497,502	904,958	296,319	237,290	83,806	23,035	9,308	3,082,749
Total mental health-related patient days	1,062,414	522,894	921,046	311,286	240,797	86,942	23,181	9,411	3,177,971

(a) Includes all separations with a mental health principal diagnosis as defined in Appendix 1 and all separations for which specialised psychiatric care was reported.

(b) Count of patient days for separations with some specialised psychiatric care.

Note: ICD-9-CM data reported by Queensland, Western Australia, South Australia and Tasmania have been mapped to ICD-10-AM. Comparisons with data reported by other jurisdictions in ICD-10-AM should be made cautiously, with reference to the classifications and maps, as appropriate.

Table 2.3: Summary of mental health-related separations, by principal diagnosis in ICD-10-AM groupings^(a) and type of care, Australia, 1998-99

Principal diagnosis	Separations with specialised psychiatric care						Other mental health-related separations			Total patient days
	Same day separations	Overnight separations	Total separations	Total patient days	Total psychiatric care days ^(b)	Total separations	Same day separations	Overnight separations	Total separations	
F00–F09 Organic, including symptomatic, mental disorders	113	1,663	1,776	108,996	88,273	208	4,871	5,079	159,463	
F00–F03 Dementia	242	1,301	1,543	95,732	89,377	549	6,158	6,707	87,648	
F04–F09 Other organic mental disorders	355	2,964	3,319	204,728	177,650	757	11,029	11,786	247,111	
F10–F19 Mental and behavioral disorders due to psychoactive substance use	2,862	4,359	7,221	126,703	107,680	4,993	11,276	16,269	70,482	
F10 Mental and behavioural disorders due to use of alcohol	1,662	5,621	7,283	48,430	44,822	1,296	6,323	7,619	33,352	
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	4,524	9,980	14,504	175,133	152,502	6,289	17,599	23,888	103,834	
F20–F29 Schizophrenia, schizotypal and delusional disorders	4,182	18,311	22,493	871,557	692,839	1,190	1,777	2,967	20,148	
F20 Schizophrenia	2,869	9,021	11,890	186,012	162,800	1,177	1,469	2,646	14,484	
F21–F29 Other schizophrenic, schizotypal, delusional disorders	7,057	27,332	34,383	1,057,569	855,639	2,367	3,246	5,613	34,632	
F30–F39 Mood (affective) disorders	41	797	838	29,141	27,957	110	234	344	2,393	
F30 Manic episode	3,403	9,030	12,433	211,069	191,980	822	1,208	2,030	13,553	
F31 Bipolar affective disorders	24,041	19,181	43,222	369,308	328,939	5,957	9,503	15,460	91,382	
F32–F33 Depressive disorders	2,785	1,922	4,707	26,430	21,910	457	1,456	1,913	11,642	
F34–F39 Other mood (affective) disorders	30,270	30,930	61,200	635,948	570,786	7,346	12,401	19,747	118,970	
F40–F48 Neurotic, stress-related and somatoform disorders	13,436	11,830	25,266	124,212	104,562	5,140	9,531	14,671	60,586	
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	4,808	1,047	5,855	34,828	31,647	295	757	1,052	14,916	
F50 Eating disorders	159	399	558	5,523	5,381	266	2,633	2,899	11,231	
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	4,967	1,446	6,413	40,351	37,028	561	3,390	3,951	26,147	
F60–F69 Disorders of adult personality and behaviour	2,521	4,913	7,434	48,677	44,563	319	872	1,191	5,810	
F70–F79 Mental retardation	25	161	186	44,084	43,807	73	84	157	1,985	
F80–F89 Disorders of psychological development	221	164	385	41,405	27,292	532	182	714	3,671	
F90–F98 Disorders onset usually occurring in childhood, adolescence	2,422	804	3,226	12,839	9,574	232	661	893	3,427	
F99 Mental disorder not otherwise specified	14	104	118	894	892	29	19	48	85	
Z03.2 Observation for suspected mental and behavioural disorder	15	110	125	866	793	12	36	48	89	
Other reasons for encounter associated with mental disorders ^(c)	414	108	522	5,893	5,746	51	91	142	471	
Disorders not related to mental health ^(a)	5,257	5,433	10,690	147,835	112,995	0	0	0	0	
Not reported	22	786	808	30,720	30,720					
Total	71,514	97,065	168,579	2,571,154	2,174,549	23,708	59,141	82,849	606,818	

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Disorders not related to mental health'.

(b) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

(c) Including Z09.3, Z13.4, Z13.5, Z24.3, Z76.5, Z91.4 and Z91.5 only.

Table 2.4: All separations in Major Diagnostic Categories 19 and 20 or with specialised psychiatric care, by AR-DRG and type of care, Australia, 1998–99

AR-DRG Description ^(a)	Separations with specialised psychiatric care						Separations with no specialised psychiatric care		
	Same day separations			Total patient days			Total psychiatric care days ^(b)		Total patient days
	Same day separations	Overnight separations	Total separations	Total patient days	Total psychiatric care days	separations	Same day separations	Overnight separations	Total separations
U40Z Mental health treatment, same day with Ect	5,185	0	5,185	5,185	5,102	2,177	0	2,177	2,177
U60Z Mental health treatment, same day without Ect	56,240	0	56,240	56,240	44,042	13,159	0	13,159	13,159
U61A Schizophrenia disorders with involuntary Mhls	0	12,441	12,441	619,238	486,296	0	13	13	96
U61B Schizophrenia disorders without involuntary Mhls	0	11,182	11,182	377,520	315,744	0	1,618	1,618	18,551
U62A Paranoid & acute psych disorder with Csc/c/involuntary Mhls	0	2,100	2,100	31,758	28,128	0	122	122	2,092
U62B Paranoid & acute psych disorder without Csc/c/involuntary Mhls	0	1,591	1,591	21,599	18,594	0	615	615	4,752
U63A Major affective disorder with Csc/c or Age > 69 without Csc/c	0	4,674	4,674	145,774	128,202	0	970	970	15,466
U63B Major affective disorder Age < 70 without Csc/c	0	21,367	21,367	397,976	364,366	0	3,284	3,284	29,426
U64Z Other affective & somatoform disorders	0	5,688	5,688	69,349	60,816	0	4,814	4,814	34,010
U65Z Anxiety disorders	0	1,918	1,918	25,653	21,226	0	8,443	8,443	31,991
U66Z Eating & obsessive-compulsive disorders	0	1,471	1,471	36,275	33,212	0	620	620	10,734
U67Z Personality disorders & acute reactions	0	14,594	14,594	126,112	109,724	0	4,033	4,033	25,726
U68Z Childhood mental disorders	0	433	433	45,238	29,508	0	545	545	3,409
<i>Total MDC 19</i>	<i>61,425</i>	<i>77,459</i>	<i>138,884</i>	<i>1,957,917</i>	<i>1,644,960</i>	<i>15,336</i>	<i>25,077</i>	<i>40,413</i>	<i>191,589</i>
V60Z Alcohol intoxication and withdrawal	914	1,879	2,793	46,613	31,844	2,059	3,674	5,733	18,455
V61A Drug intoxication and withdrawal with CC	343	488	831	6,345	5,954	44	245	289	1,579
V61B Drug intoxication and withdrawal without CC	270	1,994	2,264	17,302	15,516	272	676	948	3,102
V62A Alcohol use disorder & dependence	0	2,390	2,390	31,730	30,120	0	3,201	3,201	25,523
V62B Alcohol use disorder & dependence, same day	1,946	0	1,946	1,946	1,749	1,051	0	1,051	1,051
V63Z Opioid use disorder & dependence	410	1,850	2,260	11,334	10,436	259	2,491	2,750	10,826
V64Z Other drug use disorder & dependence	715	1,446	2,161	15,006	14,397	211	860	1,071	5,727
<i>Total MDC 20</i>	<i>4,598</i>	<i>10,047</i>	<i>14,645</i>	<i>130,276</i>	<i>110,016</i>	<i>3,896</i>	<i>11,147</i>	<i>15,043</i>	<i>66,263</i>
<i>All other MDCs</i>	<i>5,491</i>	<i>9,559</i>	<i>15,050</i>	<i>482,961</i>	<i>419,575</i>				
Total	71,514	97,065	168,579	2,571,154	2,174,551	19,232	36,224	55,456	257,852

(a) Refer to Appendix 2 for full AR-DRG category names.

(b) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

Abbreviations

- Ect Electroconvulsive therapy
- Mhls Mental health legal status
- Csc/c Catastrophic/severe complications and comorbidities
- Psych Psychotic
- CC Complications and comorbidities

Table 2.5: Separations, patient days and psychiatric care days for separations with specialised psychiatric care, by hospital type, States and Territories, 1998–99

	NSW	Vic	QLD	WA ^(a)	SA	Tas	ACT	NT	Total ^(b)
Public acute									
Same day separations	10,070	3,258	4,947	3,761	1,970	803	36	38	24,883
Overnight separations	15,293	17,082	15,398	5,146	3,943	1,845	741	724	60,172
<i>Total separations</i>	<i>25,363</i>	<i>20,340</i>	<i>20,345</i>	<i>8,907</i>	<i>5,913</i>	<i>2,648</i>	<i>777</i>	<i>762</i>	<i>85,055</i>
Total patient days	265,109	278,523	189,437	103,283	55,120	16,404	8,677	8,033	924,586
Total psychiatric care days	251,974	278,523	182,888	n.a.	55,120	16,404	8,677	7,419	801,007
Private									
Same day separations	11,821	15,850	6,953	9,731	12	n.a.	0	n.a.	44,367
Overnight separations	5,909	5,422	3,519	2,444	2,487	n.a.	50	n.a.	19,831
<i>Total separations</i>	<i>17,730</i>	<i>21,272</i>	<i>10,472</i>	<i>12,175</i>	<i>2,499</i>	<i>n.a.</i>	<i>50</i>	<i>n.a.</i>	<i>64,198</i>
Total patient days	117,892	105,493	79,056	41,501	42,247	n.a.	868	n.a.	387,057
Total psychiatric care days	115,399	105,493	78,107	n.a.	42,247	n.a.	868	n.a.	342,114
Public psychiatric									
Same day separations	2,011	2	23	53	167	8	2,264
Overnight separations	8,224	568	1,463	2,712	3,571	524	17,062
<i>Total separations</i>	<i>10,235</i>	<i>570</i>	<i>1,486</i>	<i>2,765</i>	<i>3,738</i>	<i>532</i>	<i>..</i>	<i>..</i>	<i>19,326</i>
Total patient days	471,239	15,423	512,878	114,656	98,655	46,660	1,259,511
Total psychiatric care days	243,160	15,423	512,878	114,656	98,655	46,660	1,031,432
All hospitals									
Same day separations	23,902	19,110	11,923	13,545	2,149	811	36	38	71,514
Overnight separations	29,426	23,072	20,380	10,302	10,001	2,369	791	724	97,065
<i>Total separations</i>	<i>53,328</i>	<i>42,182</i>	<i>32,303</i>	<i>23,847</i>	<i>12,150</i>	<i>3,180</i>	<i>827</i>	<i>762</i>	<i>168,579</i>
Total patient days	854,240	399,439	781,371	259,440	196,022	63,064	9,545	8,033	2,571,154
Total psychiatric care days	610,533	399,439	773,873	114,656	196,022	63,064	9,545	7,419	2,174,551

(a) Psychiatric care days from WA were not available for public acute and private hospitals.

(b) Australian total does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

.. Not applicable

n.a. Not available

Table 2.6: Separations with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings,^(a) Australia, 1998–99

Principal diagnosis	Separations		Seps per 1,000 pop'n ^(b)		Total patient days	Total psychiatric care days ^(c)	Average length of stay per separation (overnight)	Psychiatric days per care days	Patient days per 1,000 pop'n ^(b)	Psych days per care days
	Same day	Overnight	Total	patient days	psychiatric care days ^(c)					
F00–F09 Organic, including symptomatic, mental disorders	113	1,663	1,776	0.1	108,996	88,273	65.5	53.0	5.8	4.7
F00–F03 Dementia	242	1,301	1,543	0.1	95,732	89,377	73.4	68.5	5.1	4.7
F04–F09 Other organic mental disorders	355	2,964	3,319	0.2	204,728	177,650	69.0	59.8	10.9	9.4
<i>Total</i>										
F10–F19 Mental and behavioral disorders due to psychoactive substances	2,862	4,359	7,221	0.4	126,703	107,680	28.4	24.0	6.7	5.7
F10 Mental and behavioural disorders due to use of alcohol										
F11 Mental and behavioural disorders due to other psychoactive substances use	1,662	5,621	7,283	0.4	48,430	44,822	8.3	7.7	2.6	2.4
<i>Total</i>	4,524	9,980	14,504	0.8	175,133	152,502	17.1	14.8	9.3	8.1
F20–F29 Schizophrenia, schizotypal and delusional disorders	4,182	18,311	22,493	1.2	871,557	692,839	47.4	37.6	46.3	36.8
F20 Schizophrenia	2,869	9,021	11,890	0.6	186,012	162,800	20.3	17.7	9.9	8.6
F21–F29 Other schizophrenic, schizotypal, delusional disorders	7,051	27,332	34,383	1.8	1,057,569	855,639	38.4	31.0	56.1	45.4
<i>Total</i>										
F30–F39 Mood (affective) disorders	41	797	838	0.0	29,141	27,957	36.5	35.0	1.5	1.5
F30 Manic episode	3,403	9,030	12,433	0.7	211,069	191,980	23.0	20.9	11.2	10.2
F31 Bipolar affective disorders	24,041	19,181	43,222	2.3	369,308	328,939	18.0	15.9	19.6	17.5
F32–F33 Depressive disorders	2,785	1,922	4,707	0.2	26,430	21,910	12.3	10.0	1.4	1.2
F34–F39 Other mood (affective) disorders	30,270	30,930	61,200	3.2	635,948	570,786	19.6	17.5	33.8	30.3
<i>Total</i>										
F40–F48 Neurotic, stress-related and somatoform disorders	13,436	11,830	25,266	1.3	124,212	104,562	9.4	7.7	6.6	5.5
F50–F59 Behavioural syndromes associated with physiological factors	4,808	1,047	5,855	0.3	34,828	31,647	28.7	25.6	1.8	1.7
F50 Eating disorders										
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	159	399	558	0.0	5,523	5,381	13.4	13.1	0.3	0.3
<i>Total</i>	4,967	1,446	6,413	0.3	40,351	37,028	24.5	22.2	2.1	2.0
F60–F69 Disorders of adult personality and behaviour	2,521	4,913	7,434	0.4	48,677	44,563	9.4	8.6	2.6	2.4
F70–F79 Mental retardation	25	161	186	0.0	44,084	43,807	273.7	271.9	2.3	2.3
F80–F89 Disorders of psychological development	221	164	385	0.0	41,405	27,292	251.1	165.1	2.2	1.4
F90–F98 Disorders onset usually occurring in childhood, adolescence	2,422	804	3,226	0.2	12,839	9,574	13.0	8.9	0.7	0.5
F99 Mental disorder not otherwise specified	14	104	118	0.0	894	892	8.5	8.4	0.0	0.0
Z03–Z03 Observation for suspected mental and behavioural problems	15	110	125	0.0	866	793	7.7	7.1	0.0	0.0
Other Z codes associated with mental disorders ^(d)	414	108	522	0.0	5,893	5,746	50.7	49.4	0.3	0.3
Other principal diagnoses not related to mental health ^(a)	5,257	5,433	10,690	0.6	147,835	112,997	26.2	19.8	7.8	6.0
Not reported	22	786	808	0.0	30,720	30,720	39.1	39.1	1.6	1.6
Total	71,514	97,065	168,579	8.9	2,571,154	2,174,551	25.8	21.7	136.5	115.4

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses' not related to mental health.¹

(b) Population rates are crude rates based on 31 December 1998 estimated resident population data.

(c) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

(d) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5 only.

Table 2.7: Separations with specialised psychiatric care by principal diagnosis in ICD-10-AM chapter groupings, Australia, 1998–99

Principal diagnosis	Same day separations			Overnight separations			Total separation population ^(a)			Separations per 1,000 patient days			Total psychiatric care days ^(b) (overnight)			Psychiatric care days per separation			Total patient days per 1,000 population ^(a)			Total psychiatric care days per 1,000 population			
	2	30	32	10	49	59	< 0.01	761	450	25.3	14.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
A00–B99 Certain infectious and parasitic diseases	2	30	32	10	49	59	< 0.01	761	450	25.3	14.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
C00–D48 Neoplasms																									
D50–D89 Diseases of the blood and blood-forming organs	3	9	12	< 0.01	175	155	19.1	< 0.01	175	14.8	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
E00–E90 Endocrine, nutritional and metabolic diseases	8	59	67	< 0.01	1,200	879	20.2	65,806	156,434	8.3	2,385,840	2,024,295	25.6	21.6	126.6	107.4	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	
F00–F99 Mental and behavioural disorders	789	90,628	1,607	0.1	50,831	43,133	61.2	81.8	1,607	51.8	61.2	51.8	2.7	2.3	< 0.01	< 0.01	0.0	0.0	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	
G00–G99 Diseases of the nervous system	18	13	31	< 0.01	172	71	11.8	52	53	53	61	61	9.0	9.0	9.0	9.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	
H00–H59 Diseases of the eye and adnexa	3	74	77	< 0.01	8,805	8,220	118.9	74	77	69	859	575	19.4	12.8	0.0	0.0	0.5	0.4	0.0	0.0	0.0	0.0	0.0	0.0	
H60–H95 Diseases of the ear and mastoid process	26	43	69	< 0.01	990	605	13.4	26	43	100	990	605	13.4	8.0	8.0	8.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	
I00–I99 Diseases of the circulatory system	28	72	100	< 0.01	307	244	15.7	28	28	< 0.01	307	244	15.7	12.4	12.4	12.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
J00–J99 Diseases of the respiratory system	9	19	28	< 0.01	1,626	1,293	12.2	9	19	571	0.0	1,293	12.2	8.7	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
K00–K93 Diseases of the digestive system	477	94	571	0.0	702	429	15.6	48	42	90	< 0.01	702	429	15.6	9.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
L00–L99 Diseases of the skin and subcutaneous tissue	488	410	898	0.0	5,404	4,111	12.0	488	410	898	0.0	5,404	4,111	12.0	8.8	0.3	0.3	0.2	0.2	0.0	0.0	0.0	0.0		
M00–M99 Diseases of the musculoskeletal system and connective tissue	477	94	571	0.0	1,626	1,293	12.2	477	94	571	0.0	1,626	1,293	12.2	8.7	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0		
N00–N99 Diseases of the genito-urinary system	48	42	90	< 0.01	702	429	15.6	488	410	898	0.0	5,404	4,111	12.0	8.8	0.3	0.3	0.2	0.2	0.0	0.0	0.0	0.0		
O00–O99 Pregnancy, childbirth and the puerperium																									
P00–P96 Certain conditions originating in the perinatal period	2	1	3	< 0.01	26	22	24.0	701	379	1,080	0.1	4,350	2,525	9.6	4.8	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1		
Q00–Q99 Congenital malformations, deformations, and chromosomal abnormalities	1	14	15	< 0.01	5,945	2,361	424.6	337	2,160	2,497	0.1	17,755	14,423	8.1	6.5	0.9	0.8	0.8	0.8	0.8	0.8	0.8	0.8		
R00–R99 Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	2,684	1,364	4,048	0.2	53,605	39,258	37.3	2,684	1,364	4,048	0.2	53,605	39,258	37.3	26.8	2.8	2.1	2.1	2.1	2.1	2.1	2.1	2.1		
S00–T98 Injury, poisoning and certain other consequences of external causes	22	786	808	0.0	30,720	30,720	39.1	Not reported	22	786	808	0.0	30,720	30,720	39.1	39.1	1.6	1.6	1.6	1.6	1.6	1.6	1.6		
Total	71,514	97,065	168,579	9.0	2,571,154	2,174,551	25.8	71,514	97,065	168,579	9.0	2,571,154	2,174,551	25.8	21.7	136.5	115.4								

(a) Population rates are crude rates based on the 31 December 1998 estimated resident population data.

(b) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

Table 2.8: Separations, patient days and psychiatric care days of separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM groupings^(a) and hospital type, Australia, 1998–99

Principal diagnosis	Public acute hospitals						Private hospitals					
	Separations			Total patient days			Separations			Total patient days		
	Same day	Overnight	Total	psychiatric care days ^(b)	Same day	Total	Overnight	Total	psychiatric care days ^(b)	Same day	Total	psychiatric care days ^(b)
F00–F09 Organic, including symptomatic, mental disorders	105	974	1,079	38,406	18,004	7	160	167	3,824	3,687	3,824	3,687
F00–F03 Dementia	174	812	986	20,417	14,632	62	193	255	3,672	3,165	3,672	3,165
Other organic mental disorders	279	1,786	2,065	58,823	32,636	69	353	422	7,496	6,852	7,496	6,852
Total												
F10–F19 Mental and behavioural disorders due to psychoactive substances	535	1,490	2,025	11,496	9,471	2,223	1,602	3,825	24,044	21,966		
F10 Mental and behavioural disorders due to use of alcohol	310	2,487	2,797	18,380	16,678	867	1,238	2,105	14,674	13,110		
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	845	3,977	4,822	29,876	26,149	3,090	2,840	5,930	38,718	35,076		
Total												
F20–F29 Schizophrenia, schizotypal and delusional disorders	1,589	13,523	15,112	271,150	252,049	2,128	962	3,090	19,536	17,718		
F20 Schizophrenia	1,183	6,507	7,690	106,564	95,886	1,526	904	2,430	17,730	16,101		
F21–F29 Other schizotypic, schizotypal, delusional	2,772	20,030	22,802	377,714	347,935	3,654	1,866	5,520	37,266	33,879		
Total												
F30–F39 Mood (affective) disorders	33	559	592	7,665	6,676	7	86	93	1,465	1,270		
F30 Manic episode	1,102	5,879	6,981	101,191	91,018	2,095	1,531	3,626	29,730	26,783		
F31 Bipolar affective disorders	6,626	10,234	16,880	164,966	144,379	17,141	7,452	24,593	158,387	141,583		
F32–F33 Depressive disorders	1,031	1,228	2,259	11,851	8,674	1,745	507	2,256	11,493	10,182		
F34–F39 Other mood (affective) disorders	8,792	17,900	26,692	285,673	250,747	20,988	9,576	30,564	201,075	179,878		
Total												
F40–F48 Neurotic, stress-related and somatoform disorders	3,954	7,238	11,192	53,736	42,945	9,115	2,977	12,092	55,627	47,638		
F50–F59 Behavioural syndromes associated with physiological factors	1,719	581	2,300	15,799	14,288	3,089	438	3,527	18,139	16,621		
F50 Eating disorders												
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	44	191	235	2,455	2,342	110	105	215	1,976	1,947		
Total												
F60–F69 Disorders of adult personality and behaviour	1,377	3,310	4,687	24,356	21,558	1,060	467	1,527	8,329	7,205		
F60–F69 Disorders of adult personality and behaviour	19	91	110	1,026	942	0	7	7		101		
F70–F79 Mental retardation	201	104	305	2,228	1,061	16	13	29		296		
F80–F89 Disorders of psychological development	2,303	676	2,979	10,825	8,249	57	63	120		770		
F90–F98 Disorders onset usually occurring in childhood, mental disorder not otherwise specified	13	101	114	872	870	1	3	4		22		
F99 Observation for suspected mental and behavioural disorders ^(c)	13	73	86	540	467	0	3	3		16		
Z03.2 Other Z codes associated with mental disorders ^(c)	16	74	90	1,138	1,016	394	11	405	4,686	4,661		
Other principal diagnoses not related to mental health ^(a)	2,525	3,814	6,339	54,137	44,412	2,715	1,102	3,817	12,329	7,259		
Not reported	11	226	237	5,388	5,388	9	7	16	211	211		
Total	24,883	60,172	85,055	924,586	801,005	44,367	19,831	64,198	387,057	342,114		

(continued)

Table 2.8 (continued): Separations, patient days and psychiatric care days of separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM groupings^(a) and hospital type, Australia, 1998–99

Principal diagnosis	Public psychiatric hospitals					
	Separations		Total		Total	
	Same day	Overnight	Total patient days	psychiatric care days ^(b)	patient days	psychiatric care days ^(b)
F00–F09 Organic, including symptomatic, mental disorders	1	529	530	66,766	66,582	
F00–F03 Dementia	6	296	302	71,643	71,580	
F04–F09 Other organic mental disorders	7	825	832	138,409	138,162	
<i>Total</i>						
F10–F19 Mental and behavioural disorders due to psychoactive substance use	104	1,267	1,371	91,163	76,243	
F10 Mental and behavioural disorders due to use of psychoactive substances	485	1,896	2,381	15,376	15,034	
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	589	3,163	3,752	106,539	91,277	
<i>Total</i>						
F20–F29 Schizophrenia, schizotypal and delusional disorders	465	3,826	4,291	580,871	423,072	
F20 Schizophrenia	160	1,610	1,770	61,718	50,813	
F21–F29 Other schizophrenic, schizotypal, delusional disorders	625	5,436	6,061	642,589	473,885	
<i>Total</i>						
F30–F39 Mood (affective) disorders	1	152	153	20,011	20,011	
F30 Manic episode	206	1,620	1,826	80,148	74,179	
F31 Bipolar affective disorders	274	1,495	1,769	45,955	42,977	
F32–F33 Depressive disorders	9	187	196	3,086	3,054	
F34–F39 Other mood (affective) disorders	490	3,454	3,944	149,200	140,221	
<i>Total</i>						
F40–F48 Neurotic, stress-related and somatoform disorders	367	1,615	1,982	14,849	13,979	
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	0	28	28	890	738	
F50 Eating disorders						
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	5	103	108	1,092	1,092	
<i>Total</i>						
F60–F69 Disorders of adult personality and behaviour	84	1,136	1,220	15,992	15,800	
F60 Disorders of adult personality and behaviour	5	131	136	1,982	1,830	
F70–F79 Mental retardation	6	63	69	42,957	42,764	
F80–F89 Disorders of psychological development	4	47	51	38,881	25,935	
F90–F98 Disorders onset usually occurring in childhood,	62	65	127	1,244	753	
F99 Mental disorder not otherwise specified	0	0	0	0	0	
Z032 Observation for suspected mental and behavioural disorders ^(c)	2	34	36	310	310	
Other Z codes associated with mental disorders ^(c)	4	23	27	69	69	
Other principal diagnoses not related to mental health ^(a)	17	517	534	81,369	61,326	
Not reported	2	553	555	25,121	25,121	
<i>Total</i>						
	2,264	17,062	19,326	1,259,511	1,031,432	

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Does not include psychiatric care days from WA public acute or private hospitals. The totals for these hospital types are therefore underestimates of total psychiatric care days.

(c) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4, Z91.5.

Table 2.9: Same day separations with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings,^(a) States and Territories, 1998–99

Principal diagnosis	NSW	Vic	Qld	WA ^(b)	SA	Tas	ACT	NT	Total
F00–F09 Organic, including symptomatic, mental disorders	90	21	1	0	1	0	0	0	113
F00–F03 Dementia	163	25	41	8	2	3	0	0	242
F04–F09 Other organic mental disorders	253	46	42	8	3	3	0	0	355
<i>Total</i>									
F10–F19 Mental and behavioural disorders due to psychoactive substance use	1,759	527	260	276	22	13	2	3	2,862
F10 Mental and behavioural disorders due to use of alcohol	1,169	240	168	62	13	4	5	1	1,662
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	2,928	767	428	338	35	17	7	4	4,524
<i>Total</i>									
F20–F29 Schizophrenia, schizotypal and delusional disorders	1,472	1,158	877	133	447	85	4	6	4,182
F20 Schizophrenia	1,019	702	553	420	122	46	4	3	2,869
F21–F29 Other schizophrenic, schizotypal, delusional disorders	2,491	1,860	1,430	553	569	131	8	9	7,051
<i>Total</i>									
F30–F39 Mood (affective) disorders	15	9	16	1	0	0	0	0	41
F30 Manic episode	922	1,170	425	551	173	161	1	0	3,403
F31 Bipolar affective disorders	6,342	8,525	3,915	4,465	427	360	4	3	24,041
F32–F33 Depressive disorders	365	589	642	1,154	14	20	1	0	2,785
F34–F39 Other mood (affective) disorders	7,644	10,293	4,998	6,171	614	541	6	3	30,270
<i>Total</i>									
F40–F48 Neurotic, stress-related and somatoform disorders	4,449	2,153	2,426	3,798	559	34	6	11	13,436
<i>Total</i>									
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	1,043	2,687	590	447	41	0	0	0	4,808
F50 Eating disorders	64	84	7	4	0	0	0	0	159
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	1,107	2,771	597	451	41	0	0	0	4,967
<i>Total</i>									
F60–F69 Disorders of adult personality and behaviour	776	486	257	673	308	15	3	3	2,521
F70–F79 Mental retardation	5	3	16	1	0	0	0	0	25
F80–F89 Disorders of psychological development	119	15	87	0	0	0	0	0	221
F90–F98 Disorders onset usually occurring in childhood, adolescence	2,271	42	59	40	8	0	0	2	2,422
F99 Mental disorder, not otherwise specified	2	12	0	0	0	0	0	0	14
Z03.2 Observation for suspected mental and behavioural disorder	2	1	6	3	0	1	0	2	15
Other Z codes associated with mental disorders ^(c)	7	1	403	0	0	3	0	0	414
Other principal diagnoses not related to mental health ^(a)	1,848	648	1,174	1,509	12	56	6	4	5,257
Not reported	0	12	0	0	0	10	0	0	22
<i>Total</i>									
Age-standardised rate^(d)	3.75	4.06	3.44	7.38	1.43	1.73	0.12	0.21	3.80

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category Other principal diagnoses not related to mental health.

(b) WA figures include same day group therapy sessions at some units.

(c) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

(d) Indirect age-standardisation on the estimated resident population as at 31 December 1998.

Note: ICD-9-CM data reported by Queensland, Western Australia and Tasmania have been mapped to ICD-10-AM. Comparisons with data reported by other jurisdictions in ICD-10-AM should be made cautiously, with reference to the classifications and maps, as appropriate.

Table 2.10: Overnight separations with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings,^(a) States and Territories, 1998–99

Principal diagnosis	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Total
F00–F09 Organic, including symptomatic, mental disorders	242	422	173	475	338	10	1	2	1,663
F00–F03 Dementia	302	298	289	269	113	10	11	9	1,301
F04–F09 Other organic mental disorders	544	720	462	744	451	20	12	11	2,964
<i>Total</i>									
F10–F19 Mental and behavioural disorders due to psychoactive substance use	2,361	593	551	368	382	46	19	39	4,359
F10 Mental and behavioural disorders due to use of alcohol	2,891	934	895	390	337	59	47	68	5,621
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	5,252	1,527	1,446	758	719	105	66	107	9,980
<i>Total</i>									
F20–F29 Schizophrenia, schizotypal and delusional disorders	5,686	5,035	3,999	1,439	1,434	365	161	192	18,311
F20 Schizophrenia	2,749	2,143	1,843	804	1,110	94	83	90	9,021
F21–F29 Other schizophrenic, schizotypal, delusional disorders	8,435	7,178	5,842	2,243	2,544	560	255	275	27,332
<i>Total</i>									
F30–F39 Mood (affective) disorders	239	180	153	114	73	26	7	5	797
F30 Manic episode	2,535	2,081	1,879	879	1,192	296	109	59	9,030
F31 Bipolar affective disorders	5,131	5,568	4,030	1,783	2,161	330	92	86	19,181
F32–F33 Depressive disorders	381	300	560	258	302	101	16	4	1,922
F34–F39 Other mood (affective) disorders	8,286	8,129	6,622	3,034	3,728	753	224	154	30,930
<i>Total</i>									
F40–F48 Neurotic, stress-related and somatoform disorders	2,953	2,222	2,915	1,778	1,506	273	76	107	11,830
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	308	252	291	66	116	8	4	2	1,047
F50 Eating disorders	118	123	36	75	19	23	2	3	399
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	426	375	327	141	135	31	6	5	1,446
<i>Total</i>									
F60–F69 Disorders of adult personality and behaviour	1,317	1,242	937	569	557	182	91	18	4,913
F60–F69 Disorders of adult personality and behaviour	47	13	57	14	15	5	4	6	161
F70–F79 Mental retardation	74	24	32	19	12	1	2	0	164
F80–F89 Disorders of psychological development	171	173	226	144	69	18	1	2	804
F90–F98 Disorders onset usually occurring in childhood, adolescence	15	88	0	0	0	0	1	0	104
F99 Mental disorder, not otherwise specified	30	10	21	38	3	3	1	4	110
Z03.2 Observation for suspected mental and behavioural disorder	90	1	8	2	1	3	0	3	108
Other Z codes associated with mental disorders ^(b)	1,786	762	1,485	818	261	238	52	32	5,434
Other principal diagnoses not related to mental health ^(a)	0	608	0	0	0	177	0	0	785
Not reported									
<i>Total</i>									
Age-standardised rate^(c)	4.63	4.89	5.89	5.58	6.73	5.16	2.51	3.75	5.15

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

(c) Indirect age-standardisation on the estimated resident population as at 31 December 1998.

Note: ICD-9-CM data reported by Queensland, Western Australia and Tasmania have been mapped to ICD-10-AM. Comparisons with data reported by other jurisdictions in ICD-10-AM should be made cautiously, with reference to the classifications and maps, as appropriate.

Table 2.11: Patient days for overnight separations with specialised care by principal diagnosis in ICD-10-AM groupings,^(a) States and Territories, 1998–99

Principal diagnosis	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Total
F00–F09 Organic, including symptomatic, mental disorders	12,914	13,084	22,376	38,406	21,886	157	13	47	108,883
Dementia	9,986	8,061	62,052	11,274	3,279	92	149	597	95,490
Other organic mental disorders	22,900	21,145	84,428	49,680	25,165	249	162	644	204,373
<i>Total</i>									
F10–F19 Mental and behavioural disorders due to psychoactive substance use	55,529	5,614	51,589	4,704	5,955	201	91	158	123,841
Mental and behavioural disorders due to use of alcohol	23,288	6,763	7,466	4,770	3,178	492	279	532	46,768
Mental and behavioural disorders due to other psychoactive substances use	78,817	12,377	59,055	9,474	9,133	693	370	690	170,609
<i>Total</i>									
F20–F29 Schizophrenia, schizotypal and delusional disorders	354,191	104,266	274,617	63,168	44,166	21,936	2,736	2,295	867,375
Schizophrenia	67,018	35,533	32,617	21,037	21,068	2,871	1,521	1,478	183,143
Other schizophrenic, schizotypal, delusional disorders	421,209	139,799	307,234	84,205	65,234	24,807	4,257	3,773	1,050,518
<i>Total</i>									
F30–F39 Mood (affective) disorders	3,758	2,366	15,572	5,417	1,535	314	93	45	29,100
Manic episode	58,924	36,185	61,732	18,822	21,582	7,953	1,607	861	207,666
Bipolar affective disorders	101,198	92,234	70,666	34,038	42,067	2,889	1,213	962	345,267
Depressive disorders	3,858	4,277	6,705	3,877	3,516	1,225	171	16	23,645
Other mood (affective) disorders	167,738	135,062	154,675	62,154	68,700	12,381	3,084	1,884	605,678
<i>Total</i>									
F40–F48 Neurotic, stress-related and somatoform disorders	29,730	22,088	27,106	16,624	12,882	1,483	482	381	110,776
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	10,604	7,703	6,677	1,691	3,003	164	147	31	30,020
Eating disorders									
Other behavioural syndromes associated with physiological disturbances, physical factors	1,820	1,982	303	743	119	339	4	54	5,364
<i>Total</i>	12,424	9,685	6,980	2,434	3,122	503	151	85	35,384
F60–F69 Disorders of adult personality and behaviour	12,003	9,867	9,287	7,429	4,935	1,907	514	214	46,156
Mental retardation	2,706	90	38,807	1,420	120	876	18	22	44,059
Disorders of psychological development	15,514	337	24,024	1,167	95	1	46	0	41,184
Disorders onset usually occurring in childhood, adolescence	1,869	3,188	2,052	2,748	469	85	1	5	10,417
Mental disorder not otherwise specified	73	804	0	0	0	0	3	0	880
Z03.2 Observation for suspected mental and behavioural disorder	257	114	147	308	9	5	3	8	851
Other Z codes associated with mental disorders ^(b)	1,204	10	4,219	24	4	13	0	5	5,479
Other principal diagnoses not related to mental health ^(a)	63,894	12,527	51,434	8,228	4,005	1,789	418	284	142,579
Not reported	0	13,236	0	0	0	17,461	0	0	30,697
<i>Total</i>	830,338	380,329	769,448	245,895	193,873	62,253	9,509	7,995	2,499,640
Age-standardised rate^(c)	129.81	80.39	223.91	135.61	127.55	133.10	31.50	46.37	132.66

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses' not related to mental health'.

(b) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

(c) Indirect ge-standardisation on the estimated resident population as at 31 December 1998.

Note: ICD-9-CM data reported by Queensland, Western Australia, South Australia and Tasmania have been mapped to ICD-10-AM. Comparisons with data reported by other jurisdictions in ICD-10-AM should be made cautiously, with reference to the classifications and maps, as appropriate.

Table 2.12: Total psychiatric care days for overnight separations with specialised care by principal diagnosis in ICD-10-AM groupings,^(a) States and Territories, 1998–99

Principal diagnosis	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Total
F00–F09 Organic, including symptomatic, mental disorders	12,574	13,084	21,380	n.a.	21,886	157	13	47	69,141
F00–F03 Dementia	9,473	8,061	61,370	n.a.	3,279	92	149	591	83,015
F04–F09 Other organic mental disorders	22,047	21,145	82,750	n.a.	25,165	249	162	638	152,156
<i>Total</i>									
F10–F19 Mental and behavioural disorders due to psychoactive substance use	40,029	5,614	51,403	n.a.	5,955	201	91	149	103,442
F10 Mental and behavioural disorders due to use of alcohol	22,559	6,763	7,339	n.a.	3,178	492	279	517	41,127
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	62,588	12,377	58,742	n.a.	9,133	693	370	666	144,569
<i>Total</i>									
F20–F29 Schizophrenia, schizotypal and delusional disorders	192,469	104,266	274,476	n.a.	44,166	21,936	2,736	2,213	642,262
F20 Schizophrenia	55,131	35,533	32,543	n.a.	21,068	2,871	1,521	1,079	149,746
F21–F29 Other schizophrenic, schizotypal, delusional disorders	247,600	139,799	307,019	n.a.	65,234	24,807	4,257	3,292	792,008
<i>Total</i>									
F30–F39 Mood (affective) disorders	3,710	2,366	15,553	n.a.	1,535	314	93	45	23,616
F30 Manic episode	52,381	36,185	61,430	n.a.	21,582	7,953	1,607	852	181,990
F31 Bipolar affective disorders	95,271	92,234	69,756	n.a.	42,067	2,889	1,213	943	304,373
F32–F33 Depressive disorders	3,759	4,277	6,638	n.a.	3,516	1,225	171	16	19,602
F34–F39 Other mood (affective) disorders	155,121	135,062	153,377	n.a.	68,700	12,381	3,084	1,856	529,581
<i>Total</i>									
F40–F48 Neurotic, stress-related and somatoform disorders	27,766	22,088	26,926	n.a.	12,882	1,483	482	370	91,997
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	9,724	7,703	6,474	n.a.	3,003	164	147	31	27,246
F50 Eating disorders									
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	1,812	1,982	303	n.a.	119	339	4	53	4,612
<i>Total</i>	11,536	9,685	6,777	n.a.	3,122	503	151	84	31,858
F60–F69 Disorders of adult personality and behaviour	11,493	9,867	9,225	n.a.	4,935	1,907	514	214	38,155
F60–F69 Mental retardation	2,507	90	38,807	n.a.	120	876	18	22	42,440
F70–F79 Disorders of psychological development	2,555	337	24,024	n.a.	95	1	46	0	27,058
F80–F89 Disorders onset usually occurring in childhood, adolescence	1,374	3,188	2,051	n.a.	469	85	1	5	7,173
F99 Mental disorder not otherwise specified	71	804	0	n.a.	0	0	3	0	878
Z03.2 Observation for suspected mental and behavioural disorder	249	114	147	n.a.	9	5	3	8	535
Other Z codes associated with mental disorders ^(b)	1,081	10	4,219	n.a.	4	13	0	5	5,332
Other principal diagnoses not related to mental health ^(a)	40,645	12,527	47,886	n.a.	4,005	1,789	418	221	107,491
Not reported	0	13,236	0	n.a.	0	17,461	0	0	30,697
<i>Total</i>	586,633	380,329	761,950	n.a.	193,873	62,253	9,509	7,381 2,001,928	
Age-standardised rate^(c)	91.7	80.4	221.7	n.a.	127.6	133.2	31.5	42.7	106.2

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

(c) Indirect age-standardisation on the estimated resident population as at 31 December 1998.

Note: ICD-9-CM data reported by Queensland, Western Australia, South Australia and Tasmania have been mapped to ICD-10-AM. Comparisons with data reported by other jurisdictions in ICD-10-AM should be made cautiously, with reference to the classifications and maps, as appropriate.

Table 2.13a: Same day separations for male patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and age group, Australia, 1998-99

Principal diagnosis	17 and under	18-24	25-34	35-44	45-54	55-64	65 and over	Total
F00-F09 Organic, including symptomatic, mental disorders	0	0	0	0	0	0	0	94
F00-F03 Dementia	8	3	5	31	2	63	86	198
Other organic mental disorders	8	3	5	31	2	63	180	292
Total/								
F10-F19 Mental and behavioural disorders due to psychoactive substance use	7	76	142	321	493	201	478	1,718
F10 Mental and behavioural disorders due to use of alcohol	164	312	266	64	55	32	32	925
F11-F19 Mental and behavioural disorders due to other psychoactive substances use	171	388	408	385	548	233	510	2,643
Total/								
F20-F29 Schizophrenia, schizotypal and delusional disorders	489	413	441	454	364	128	50	2,339
F20 Schizophrenia	144	166	151	244	184	11	356	1,256
F21-F29 Other schizophrenic, schizotypal, delusional disorders	633	579	592	698	548	139	406	3,595
Total/								
F30-F39 Mood (affective) disorders	0	4	6	9	3	3	0	25
F30 Manic episode	192	82	134	97	299	111	140	1,055
F31 Bipolar affective disorders	368	458	907	1,681	1,799	857	2,365	8,435
F32-F33 Depressive disorders	75	68	299	272	169	108	432	1,423
F34-F39 Other mood (affective) disorders	635	612	1,346	2,059	2,270	1,079	2,937	10,938
Total/								
F40-F48 Neurotic, stress-related and somatoform disorders	833	342	842	901	1,966	681	1,860	7,425
F40-F49 Behavioural syndromes associated with physiological disturbances and physical factors	5	69	15	8	0	14	0	111
F50 Eating disorders	0	11	7	2	0	1	34	55
F51-F59 Other behavioural syndromes associated with physiological disturbances, physical factors	5	80	22	10	0	15	34	166
Total/								
F60-F69 Disorders of adult personality and behaviour	10	77	227	74	245	49	218	900
F70-F79 Mental retardation	1	2	6	3	1	0	0	13
F80-F89 Disorders of psychological development	116	99	3	1	0	0	0	219
F90-F98 Disorders onset usually occurring in childhood, adolescence	2,179	5	1	0	2	0	0	2,187
F99 Mental disorder, not otherwise specified	2	4	2	2	0	0	0	10
Z03.2 Observation for suspected mental and behavioural disorder	1	4	2	4	1	0	0	12
Other Z codes associated with mental disorders ^(b)	1	4	10	10	21	3	1	50
Other principal diagnoses not related to mental health ^(a)	524	149	694	711	228	176	104	2,586
Not reported	0	4	4	2	2	1	0	13
Total/								
Total	5,119	2,352	4,164	4,891	5,834	2,439	6,250	31,049

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category

(b) Other principal diagnoses not related to mental health:
including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.13b: Same day separations for female patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and age group, Australia, 1998-99

Principal diagnosis	17 and under	18-24	25-34	35-44	45-54	55-64	65 and over	Total
F00-F09 Organic, including symptomatic, mental disorders	0	0	0	0	0	2	2	19
F00-F03 Dementia	0	0	3	2	3	2	34	44
F04-F09 Other organic mental disorders	0	0	3	2	3	4	51	63
Total/	0	0	3	2	3	4	51	63
F10-F19 Mental and behavioural disorders due to psychoactive substance use	8	53	135	196	502	138	112	1,144
F10 Mental and behavioural disorders due to use of alcohol	308	122	111	97	90	4	5	737
F11-F19 Mental and behavioural disorders due to other psychoactive substances use	316	175	246	293	592	142	117	1,881
Total/	316	175	246	293	592	142	117	1,881
F20-F29 Schizophrenia, schizotypal and delusional disorders	83	159	276	422	456	266	181	1,843
F20 Schizophrenia	62	126	281	451	418	178	97	1,613
F21-F29 Other schizophrenic, schizotypal, delusional disorders	145	285	557	873	874	444	278	3,456
Total/	145	285	557	873	874	444	278	3,456
F30-F39 Mood (affective) disorders	1	0	2	8	1	1	3	16
F30 Manic episode	34	183	344	508	686	294	299	2,348
F31 Bipolar affective disorders	723	1,265	2,235	3,806	3,942	1,708	1,927	15,606
F32-F33 Depressive disorders	106	86	312	435	319	80	24	1,362
F34-F39 Other mood (affective) disorders	864	1,534	2,893	4,757	4,948	2,083	2,253	19,332
Total/	864	1,534	2,893	4,757	4,948	2,083	2,253	19,332
F40-F48 Neurotic, stress-related and somatoform disorders	424	502	1,298	1,352	1,147	504	784	6,011
F50-F59 Behavioural syndromes associated with physiological disturbances and physical factors	441	2,346	1,107	495	300	8	0	4,697
F50 Eating disorders	5	2	42	13	42	0	0	104
F51-F59 Other behavioural syndromes associated with physiological disturbances, physical factors	446	2,348	1,149	508	342	8	0	4,801
Total/	446	2,348	1,149	508	342	8	0	4,801
F60-F69 Disorders of adult personality and behaviour	35	182	446	414	303	150	91	1,621
F60-F69 Disorders of adult personality and behaviour	1	0	1	5	5	0	0	12
F70-F79 Mental retardation	0	0	1	1	0	0	0	2
F80-F89 Disorders of psychological development	206	5	1	17	5	0	1	235
F90-F98 Disorders onset usually occurring in childhood, adolescence	2	0	0	0	1	1	0	4
F99 Mental disorder not otherwise specified	0	0	2	1	0	0	0	3
Z032 Observation for suspected mental and behavioural disorder	1	13	31	56	147	70	46	364
Other Z codes associated with mental disorders ^(b)	413	258	1,044	583	176	105	92	2,671
Other principal diagnoses not related to mental health ^(a)	0	0	0	4	2	1	2	9
Not reported	0	0	0	0	0	0	0	0
Total/	2,853	5,302	7,672	8,866	8,545	3,512	3,715	40,465

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.14a: Overnight separations for male patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and age group, Australia, 1998–99

Principal diagnosis		17 and under	18–24	25–34	35–44	45–54	55–64	65 and over	Total
F00–F09	Organic, including symptomatic, mental disorders	0	2	1	9	11	64	755	842
	Dementia	11	50	152	118	84	88	264	767
	Other organic mental disorders	11	52	153	127	95	152	1,019	1,609
	<i>Total</i>								
F10–F19	Mental and behavioural disorders due to psychoactive substance use	17	223	629	796	774	340	244	3,023
	Mental and behavioural disorders due to use of alcohol	224	1,433	1,457	549	119	22	20	3,824
	Mental and behavioural disorders due to other psychoactive substances use	241	1,655	2,086	1,345	893	362	264	6,847
	<i>Total</i>								
F20–F29	Schizophrenia, schizotypal and delusional disorders	268	2,917	4,362	2,810	1,288	504	314	12,463
	Schizophrenia	234	1,206	1,479	870	457	166	179	4,591
	Other schizophrenic, schizotypal, delusional disorders	502	4,123	5,841	3,680	1,745	670	493	17,054
	<i>Total</i>								
F30–F39	Mood (affective) disorders	5	112	99	82	39	29	43	409
	Manic episode	48	528	1,040	863	621	415	330	3,845
	Bipolar affective disorders	197	649	1,228	1,420	1,466	735	1,247	6,942
	Depressive disorders	37	96	162	175	127	53	60	710
	Other mood (affective) disorders	287	1,385	2,529	2,540	2,253	1,232	1,680	11,906
	<i>Total</i>								
F40–F48	Neurotic, stress-related and somatoform disorders	431	769	1,430	1,142	1,228	352	205	5,557
	Behavioural syndromes associated with physiological disturbances and physical factors								
F50	Eating disorders	11	16	8	4	2	1	1	43
F51–F59	Other behavioural syndromes associated with physiological disturbances, physical factors	4	54	31	22	5	4	2	122
	<i>Total</i>	15	70	39	26	8	5	3	166
F60–F69	Disorders of adult personality and behaviour	63	400	659	351	131	44	31	1,679
	Mental retardation	8	15	34	15	15	4	5	96
	Disorders of psychological development	51	22	21	7	8	4	0	113
	Disorders onset usually occurring in childhood, adolescence	447	39	21	18	8	3	8	544
	Mental disorder not otherwise specified	1	11	27	15	2	4	1	61
	Z03.2 Observation for suspected mental and behavioural disorder	10	17	18	17	6	1	3	72
	Other Z codes associated with mental disorders ^(b)	5	16	38	24	2	1	2	88
	Other principal diagnoses not related to mental health ^(a)	312	289	534	422	285	148	378	2,368
	Not reported	9	114	214	109	29	28	91	594
	<i>Total</i>	2,393	8,978	13,644	9,838	6,708	3,010	4,183	48,754

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.14b: Overnight separations for female patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and age group, Australia, 1998–99

Principal diagnosis	17 and under	18–24	25–34	35–44	45–54	55–64	65 and over	Total
F00–F09 Organic, including symptomatic, mental disorders	0	0	4	2	7	42	766	821
F00	6	15	65	71	54	269	269	534
Dementia	6	15	69	73	61	96	1,035	1,355
Other organic mental disorders								
Total/	6	15	69	73	61	96	1,035	1,355
F10–F19 Mental and behavioural disorders due to psychoactive substance use	16	75	283	365	361	145	91	1,336
F10 Mental and behavioural disorders due to use of alcohol	189	571	592	298	86	34	27	1,797
F11 Mental and behavioural disorders due to other psychoactive substances use	205	646	875	663	447	179	118	3,133
Total/								
F20–F29 Schizophrenia, schizotypal and delusional disorders	156	781	1,433	1,344	1,022	531	581	5,848
F20 Schizophrenia	156	545	1,169	1,067	739	357	397	4,430
F21 Other schizophrenic, schizotypal, delusional disorders	312	1,326	2,602	2,411	1,761	888	978	10,278
Total/								
F30–F39 Mood (affective) disorders	8	36	78	85	54	34	93	388
F30 Manic episode	62	430	1,086	1,227	1,045	628	707	5,185
F31 Bipolar affective disorders	380	856	2,152	2,689	2,327	1,169	2,666	12,239
F32 Depressive disorders	150	149	270	244	171	77	151	1,212
F33 Other mood (affective) disorders	600	1,471	3,586	4,245	3,597	1,908	3,617	19,024
Total/								
F40–F48 Neurotic, stress-related and somatoform disorders	685	854	1,666	1,484	896	338	350	6,273
F40 Behavioural syndromes associated with physiological disturbances and physical factors	308	374	193	94	29	4	1	1,003
F50 Eating disorders	11	50	153	52	7	3	1	277
F51 Other behavioural syndromes associated with physiological disturbances, physical factors	319	424	346	146	36	7	2	1,280
Total/								
F60–F69 Disorders of adult personality and behaviour	179	794	1,130	760	277	60	34	3,234
F60 Disorders of adult personality and behaviour	4	8	19	9	12	10	3	65
F61 Mental retardation	17	10	13	3	5	3	0	51
F62 Disorders of psychological development	189	17	24	5	13	5	7	260
F63 Disorders onset usually occurring in childhood, adolescence	5	6	11	10	10	0	1	43
F64 Mental disorder not otherwise specified	1	14	8	5	3	0	7	38
Z032 Observation for suspected mental and behavioural disorder	3	4	4	4	2	1	2	20
Other Z codes associated with mental disorders ^(b)	348	353	873	628	300	148	415	3,065
Other principal diagnoses not related to mental health ^(a)	1	6	9	13	21	10	132	192
Not reported								
Total/	2,874	5,948	11,235	10,459	7,441	3,653	6,701	48,311

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.15a: Patient days for overnight separations for male patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and age group, Australia, 1998–99

Principal diagnosis	17 and under	18–24	25–34	35–44	45–54	55–64	65 and over	Total
F00–F09 Organic, including symptomatic, mental disorders	0	79	2	299	1,293	3,747	56,826	62,246
F00–F03 Dementia	73	600	11,973	7,864	23,028	10,667	18,715	72,920
F04–F09 Other organic mental disorders	73	679	11,975	8,163	24,321	14,414	75,541	135,166
<i>Total</i>								
F10–F19 Mental and behavioural disorders due to psychoactive substance	158	1,505	4,189	17,037	18,311	31,380	22,235	94,815
F10 Mental and behavioural disorders due to use of alcohol								
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	2,470	11,124	9,947	4,236	1,081	406	1,109	30,373
<i>Total</i>	2,628	12,629	14,136	21,273	19,392	31,786	23,344	125,188
F20–F29 Schizophrenia, schizotypal and delusional disorders	6,556	120,921	187,279	91,205	89,644	42,994	30,715	569,314
F20 Schizophrenia	4,127	20,161	25,910	14,183	8,438	19,290	7,270	99,379
F21–F29 Other schizotypic, schizotypal, delusional disorders	10,683	141,082	213,189	105,388	98,082	62,284	37,985	668,693
<i>Total</i>								
F30–F39 Mood (affective) disorders	67	1,419	1,385	843	489	3,310	2,275	9,788
F30 Manic episode	617	9,364	15,694	14,558	10,555	23,654	13,175	87,617
F31 Bipolar affective disorders	2,836	6,895	14,413	19,274	24,268	16,645	38,066	122,397
F32–F33 Depressive disorders	372	693	1,424	1,945	1,518	581	1,660	8,193
F34–F39 Other mood (affective) disorders	3,892	18,371	32,916	36,620	36,830	44,190	55,176	227,995
<i>Total</i>								
F40–F48 Neurotic, stress-related and somatoform disorders	3,424	4,470	8,462	7,500	18,059	4,553	4,155	50,623
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors								
F50 Eating disorders	157	411	304	28	113	6	39	1,058
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	8	634	256	194	34	52	10	1,188
<i>Total</i>	165	1,045	560	222	147	58	49	2,246
F60–F69 Disorders of adult personality and behaviour	415	3,238	6,356	2,473	1,941	507	1,280	16,210
F60–F69 Mental retardation	46	770	22,308	332	2,451	649	1,291	27,847
F70–F79 Disorders of psychological development	1,119	23,137	110	55	14,304	581	0	39,306
F80–F89 Disorders onset usually occurring in childhood, adolescence	5,360	195	103	91	115	50	115	6,029
F90–F98 Disorders onset not otherwise specified	5	84	225	143	22	16	3	498
F99 Mental disorder not otherwise specified	40	166	178	69	60	3	9	525
Z032 Observation for suspected mental and behavioural disorder	36	208	556	331	7	10	22	1,170
Other Z codes associated with mental disorders ^(b)	2,068	4,399	12,249	26,582	11,052	8,859	15,975	81,184
Other principal diagnoses not related to mental health ^(a)	50	2,393	3,875	1,766	958	1,796	7,369	18,207
Not reported								
<i>Total</i>	30,004	212,866	327,198	211,008	227,741	169,756	222,314	1,400,887

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category

(b) Other principal diagnoses not related to mental health.

Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.15b: Patient days for overnight separations for female patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and age group, Australia, 1998–99

Principal diagnosis	17 and under	18–24	25–34	35–44	45–54	55–64	65 and over	Total
F00–F09 Organic, including symptomatic, mental disorders	0	0	176	120	233	2,691	43,417	46,637
Dementia	77	143	7,822	2,270	839	1,159	10,260	22,570
Other organic mental disorders		77	143	7,998	2,390	1,072	3,850	53,677
Total								69,207
F10–F19 Mental and behavioural disorders due to psychoactive substance	102	508	2,430	3,806	4,019	9,671	8,490	29,026
Mental and behavioural disorders due to use of alcohol								
Mental and behavioural disorders due to other psychoactive substances use	2,087	4,529	4,838	2,701	1,323	390	527	16,395
Total	2,189	5,037	7,268	6,507	5,342	10,061	9,017	45,421
F20–F29 Schizophrenia, schizotypal and delusional disorders	6,708	35,159	45,818	52,409	83,247	29,577	45,143	298,061
Schizophrenia	2,666	9,585	18,733	17,662	14,783	7,463	12,872	83,764
Other schizophrenic, schizotypal, delusional disorders	9,374	44,744	64,551	70,071	98,030	37,040	58,015	381,825
Total								
F30–F39 Mood (affective) disorders	73	527	866	1,211	806	8,830	6,999	19,312
Manic episode	1,282	17,088	17,764	19,771	19,183	22,885	22,076	120,049
Bipolar affective disorders	5,059	11,475	29,082	40,237	38,908	25,071	73,038	222,870
Depressive disorders	1,184	1,272	2,604	2,631	2,258	1,763	3,740	15,452
Other mood (affective) disorders	7,598	30,362	50,316	63,850	61,155	58,549	105,853	377,683
Total								
F40–F48 Neurotic, stress-related and somatoform disorders	6,323	5,921	13,696	12,834	10,563	4,438	6,378	60,153
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	9,193	10,491	5,820	2,558	756	143	1	28,962
Eating disorders								
Other behavioural syndromes associated with physiological disturbances, physical factors	57	630	2,608	661	144	60	16	4,176
Total	9,250	11,121	8,428	3,219	900	203	17	33,138
F60–F69 Disorders of adult personality and behaviour	1,385	5,670	10,093	7,682	2,922	910	1,284	29,946
Mental retardation	48	2,974	861	1,096	5,435	4,918	880	16,212
Disorders of psychological development	481	141	32	32	1,146	46	0	1,878
Disorders onset usually occurring in childhood, adolescence	2,377	156	224	73	1,096	59	403	4,388
Mental disorder not otherwise specified	36	42	59	81	163	0	1	382
Observation for suspected mental and behavioural disorder	3	121	41	17	21	0	123	326
Z03.2 Other Z codes associated with mental disorders ^(b)	5	8	28	15	1,776	2,441	36	4,309
Other principal diagnoses not related to mental health ^(a)	2,435	2,980	13,161	13,472	9,645	3,442	16,259	61,394
Not reported	1	61	112	1,138	542	1,219	9,418	12,491
Total	41,582	109,481	176,868	182,477	199,808	127,176	261,361	1,098,753

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category

(b) Other principal diagnoses not related to mental health.

Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.16a: Total psychiatric care days^(a) for overnight separations for male patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(b) and age group, Australia, 1998–99

Principal diagnosis	17 and under	18–24	25–34	35–44	45–54	55–64	65 and over	Total
F00–F09 Organic, including symptomatic, mental disorders								
F00–F03 Dementia	52	79	2	185	1261	3502	45963	50,992
F04–F09 Other organic mental disorders	52	544	11533	7740	22483	9803	16878	69,033
Total	104	623	11535	7925	23744	13305	62841	120,025
F10–F19 Mental and behavioural disorders due to psychoactive substance use								
F10 Mental and behavioural disorders due to use of alcohol	156	1430	3893	16311	10853	26182	18518	77,343
F11–F19 Mental and behavioural disorders due to other psychoactive substances use	1838	10463	9422	4035	905	406	1063	28,132
Total	1994	11893	13315	20346	11758	26588	19581	105,475
F20–F29 Schizophrenia, schizotypal and delusional disorders								
F20 Schizophrenia	6225	80496	162837	80519	70779	27272	25611	453,739
Other schizophrenic, schizotypal, delusional disorders	3315	18475	24224	13050	8011	8713	6868	82,656
Total	9540	98971	187061	93569	78790	35985	32479	536,395
F30–F39 Mood (affective) disorders								
F30 Manic episode	63	1282	1311	658	439	3304	2199	9,256
F31 Bipolar affective disorders	531	8677	14732	13407	9956	22651	11149	81,033
F32–F33 Depressive disorders	2381	6181	12707	17546	21999	15314	33272	109,400
F34–F39 Other mood (affective) disorders	311	559	1230	1794	1380	532	1494	7,300
Total	3286	16699	29980	33405	33774	41801	48114	207,059
F40–F48 Neurotic, stress-related and somatoform disorders								
F40–F48 Behavioural syndromes associated with physiological disturbances and physical factors	2803	3694	7153	6432	16324	4039	3124	43,569
F50–F59 Eating disorders	107	382	222	28	113	6	39	897
F50–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	8	627	225	185	34	52	10	1,141
Total	115	1009	447	213	147	58	49	2,038
F60–F69 Disorders of adult personality and behaviour								
F60–F69 Disorders of adult personality and behaviour	390	3079	6015	2322	1545	414	882	14,647
F70–F79 Mental retardation	41	765	22295	332	2451	649	1042	27,575
F80–F89 Disorders of psychological development	964	23135	105	55	1334	581	53	26,174
F90–F98 Disorders onset usually occurring in childhood, adolescence	3706	169	102	89	115	50	53	4,284
F99 Mental disorder not otherwise specified	5	84	223	143	22	16	3	496
Z032 Observation for suspected mental and behavioural disorder	35	166	170	59	47	3	9	489
Other Z codes associated with mental disorders ^(c)	32	204	454	315	7	10	20	1,042
Other principal diagnoses not related to mental health ^(b)	1,273	4,081	11,257	15,518	10,149	8,468	12,531	63,277
Not reported	50	2,393	3,875	1,766	958	1,796	7,369	18,207
Total	24,286	166,965	293,987	182,489	181,165	133,763	188,097	1,170,752

(a) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

(b) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category. Other principal diagnoses not related to mental health.

(c) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.16b: Total psychiatric care days^(a) for overnight separations for female patients with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(b) and age group, Australia, 1998–99

Principal diagnosis	17 and under	18–24	25–34	35–44	45–54	55–64	65 and over	Total
F00–F09 Organic, including symptomatic, mental disorders								
F00–F03 Dementia	72	140	59	120	233	2,296	34,460	37,168
F04–F09 Other organic mental disorders	72	140	7,593	2,189	572	816	8,727	20,109
Total			7,652	2,309	805	3,112	43,187	57,277
F10–F19 Mental and behavioural disorders due to psychoactive substance use								
F10 Mental and behavioural disorders due to use of alcohol	83	500	2,284	3,234	3,807	9,428	8,408	27,744
Mental and behavioural disorders due to other psychoactive substances use	1,898	4,228	4,217	2,561	1,289	390	504	15,087
Total	1,981	4,728	6,501	5,795	5,096	9,818	8,912	42,831
F20–F29 Schizophrenia, schizotypal and delusional disorders								
F20 Schizophrenia	6,586	26,745	37,157	34,636	65,526	28,751	35,645	235,046
F21–F29 Other schizophrenic, schizotypal, delusional disorders	2,376	8,559	17,588	16,589	14,049	6,745	11,789	77,695
Total	8,962	35,304	54,745	51,225	79,575	35,496	47,434	312,741
F30–F39 Mood (affective) disorders								
F30 Manic episode	53	500	853	1,121	702	8,790	6,642	18,661
F31 Bipolar affective disorders	1,074	16,518	16,117	17,684	17,657	19,873	19,101	108,024
F32–F33 Depressive disorders	4,149	10,211	26,469	35,888	34,764	22,506	65,973	199,960
F34–F39 Other mood (affective) disorders	957	1,052	2,246	2,364	2,016	1,541	2,803	12,979
Total	6,233	28,281	45,685	57,057	55,139	52,710	94,519	339,624
F40–F48 Neurotic, stress-related and somatoform disorders								
F40–F48 Behavioural syndromes associated with physiological disturbances and physical factors	5,483	4,546	11,682	10,921	9,147	4,031	5,532	51,342
F50 Eating disorders								
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors	8,307	9,375	5,403	2,442	718	143	1	26,389
Total								
F60–F69 Disorders of adult personality and behaviour								
F60–F69 Disorders of adult personality and behaviour	54	619	2,587	644	144	17	16	4,081
F70–F79 Mental retardation	8,361	9,994	7,990	3,086	862	160	17	30,470
F80–F89 Disorders of psychological development	1,297	5,212	9,600	7,050	2,783	910	1,205	28,057
F90–F98 Disorders onset usually occurring in childhood, adolescence	48	2,974	856	1,096	5,435	4,918	880	16,207
F99 Mental disorder not otherwise specified	115	139	31	32	534	46	9	897
Z03.2 Observation for suspected mental and behavioural disorder	1,263	109	203	73	1,095	9	156	2,908
Other Z codes associated with mental disorders ^(c)	36	42	59	81	163	1	382	
Other principal diagnoses not related to mental health ^(b)	3	114	28	17	20	108	290	
Not reported	5	8	28	15	1,776	2,441	17	4,290
Total	1,725	2,186	7,776	12,396	4,910	2,941	14,036	45,970
Not reported	1	61	112	1,138	542	1,219	9,418	12,491
Total	35,585	93,838	152,948	152,291	117,811	225,422	945,777	

(a) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

(b) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(c) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5.

Table 2.17: Separations with specialised psychiatric care by marital status and sex, States and Territories 1998–99

Marital status	NSW ^(a)	Vic ^(b)	Qld	WA	SA ^(a)	Tas	ACT	NT	Total
Males									
Never married	12,842	8,663	8,242	5,105	2,746	853	312	355	39,118
Widowed	1,130	4,444	257	556	136	53	1	5	6,582
Divorced	1,005	1,848	788	816	289	106	28	16	4,896
Separated	719	67	773	775	281	64	19	14	2,712
Married (including de facto)	4,071	120	3,639	4,262	962	375	61	84	13,574
Not stated/inadequately described	2,498	1,202	591	147	435	17	19	37	4,946
Not reported	7,085	0	0	0	890	0	0	0	7,975
Total males	29,350	16,344	14,290	11,661	5,739	1,468	440	511	79,803
Females									
Never married	5,442	10,765	7,116	4,692	1,679	561	205	120	30,580
Widowed	880	9,567	1,154	1,020	399	238	8	5	13,271
Divorced	893	4,069	1,619	1,317	491	165	39	6	8,599
Separated	627	36	947	740	346	157	24	7	2,884
Married (including de facto)	3,036	85	6,562	4,284	1,653	566	99	96	16,381
Not stated/inadequately described	1,843	1,316	615	133	235	25	12	17	4,196
Not reported	11,257	0	0	0	1,608	0	0	0	12,865
Total females	23,978	25,838	18,013	12,186	6,411	1,712	387	251	88,776
Persons									
Never married	18,284	19,428	15,358	9,797	4,425	1,414	517	475	69,698
Widowed	2,010	14,011	1,411	1,576	535	291	9	10	19,853
Divorced	1,898	5,917	2,407	2,133	780	271	67	22	13,495
Separated	1,346	103	1,720	1,515	627	221	43	21	5,596
Married (including de facto)	7,107	205	10,201	8,546	2,615	941	160	180	29,955
Not stated/inadequately described	4,341	2,518	1,206	280	670	42	31	54	9,142
Not reported	18,342	0	0	0	2,498	0	0	0	20,840
Total persons	53,328	42,132	32,303	23,847	12,150	3,180	827	762	168,579

(a) Data not available for private hospitals for NSW and SA.

(b) Victorian data should be used with caution.

Table 2.18: Area of usual residence^(a) of patients for separations with specialised psychiatric care, by hospital type, States and Territories, 1998–99

Area of usual residence	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Total
Public acute hospitals									
Metropolitan	21,423	14,879	14,535	8,096	5,478	1,070	718	378	66,577
Rural	3,487	4,848	4,952	638	369	1,569	35	48	15,946
Remote	47	88	292	111	36	6	0	304	884
Not reported	406	525	566	62	30	3	24	32	1,648
<i>Total public acute hospitals</i>	<i>25,363</i>	<i>20,340</i>	<i>20,345</i>	<i>8,907</i>	<i>5,913</i>	<i>2,648</i>	<i>777</i>	<i>762</i>	<i>85,055</i>
Private hospitals									
Metropolitan	16,442	20,268	9,468	11,071	2,175	n.a.	17	n.a.	59,441
Rural	1,265	916	931	870	307	n.a.	29	n.a.	4,318
Remote	12	6	41	224	15	n.a.	0	n.a.	298
Not reported	11	82	32	10	2	n.a.	4	n.a.	141
<i>Total private hospitals</i>	<i>17,730</i>	<i>21,272</i>	<i>10,472</i>	<i>12,175</i>	<i>2,499</i>	<i>n.a.</i>	<i>50</i>	<i>n.a.</i>	<i>64,198</i>
Public psychiatric hospitals									
Metropolitan	7,940	490	810	2,248	2,780	397	14,665
Rural	1,434	71	541	254	743	131	3,174
Remote	76	0	34	150	54	0	314
Not reported	785	9	101	113	161	4	1,173
<i>Total public psychiatric hospitals</i>	<i>10,235</i>	<i>570</i>	<i>1,486</i>	<i>2,765</i>	<i>3,738</i>	<i>532</i>	<i>..</i>	<i>..</i>	<i>19,326</i>
All hospitals									
Metropolitan	45,805	35,637	24,813	21,415	10,433	1,467	735	378	140,683
Rural	6,186	5,835	6,424	1,762	1,419	1,700	64	48	23,438
Remote	135	94	367	485	105	6	0	304	1,496
Not reported	1,202	616	699	185	193	7	28	32	2,962
Total hospitals	53,328	42,182	32,303	23,847	12,150	3,180	827	762	168,579
Age-standardised population rate^(b)									
All separations per 1,000 population									
Metropolitan	9.09	9.56	11.56	9.38	5.89	7.24	1.35	2.06	9.98
Rural	1.23	1.57	2.99	0.77	0.80	8.39	0.12	0.26	1.66
Remote	2.40	7.07	1.86	13.13	0.57	2.17	..	3.46	2.59
Overnight separations per 1,000 population									
Metropolitan	4.49	4.72	6.46	3.76	4.76	5.14	1.27	2.00	5.18
Rural	1.00	1.21	2.50	0.50	0.73	6.39	0.12	0.25	1.33
Remote	0.03	0.02	0.16	0.13	0.06	0.03	..	1.64	0.09

(a) Defined according to the *Rural, Remote and Metropolitan Areas Classification*, 1991 Census edition. See Glossary for more information.

(b) Rates were indirectly age-standardised using the estimated resident population as at 31 December 1998.

.. Not applicable

n.a. Not available

Table 2.19: Specialised care separations reported for Aboriginal and Torres Strait Islander patients,^(a) by principal diagnosis in ICD-10-AM groupings, Australia, 1998-99

Principal diagnosis	Separations			Separations			Patient			Psychiatric		
	Same day separations	Overnight separations	Total separations	per 1,000 ATSI ^(b)	Total patient days	per 1,000 ATSI ^(b)	Total psychiatric care days	per 1,000 ATSI ^(b)	Average stay	length of care days per stay	Psychiatric overnight separation	
F00–F09 Organic, including symptomatic, mental disorders	2	63	65	0.2	4,731	16.0	4,731	16.0	75.1	75.1		
F10–F19 Mental and behavioural disorders due to psychoactive substance use	47	495	542	1.6	5,165	17.5	4,912	16.6	10.3	10.3	9.8	
F20–F29 Schizophrenia, schizotypal and delusional disorders	30	902	932	2.7	39,678	134.2	38,264	129.4	44.0	44.0	42.4	
F30–F39 Mood (affective) disorders	32	491	523	1.5	6,343	21.5	5,848	19.8	12.9	12.9	11.8	
F40–F49 Neurotic, stress-related and somatoform disorders	163	339	502	1.5	2,163	7.3	1,785	6.0	5.9	5.9	4.8	
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	7	30	37	0.1	383	1.3	383	1.3	12.5	12.5	12.5	
Other mental health disorders ^(d)	78	193	271	0.8	4,314	14.6	4,137	14.0	21.9	21.9	21.0	
Other principal diagnoses not related to mental health ^(e)	24	165	189	0.6	2,639	8.9	2,245	8.9	15.8	15.8	13.5	
Total	383	2,678	3,061	9.0	65,416	221.2	62,305	210.7	24.3	24.3	23.1	
All patients	71,514	97,065	168,579	8.9	2,571,154	136.5	2,083,835	115.4	25.8	25.8	20.7	

(a) A full definition of Aboriginal and Torres Strait Islander status is in the Glossary. The completeness of Aboriginal and Torres Strait Islander status varies, hence these data should be used with caution.

(b) Rates are indirectly age-standardised based on December 1998 Estimated Resident Population and the December 1998 Aboriginal and Torres Strait Islander population projections.

(c) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

(d) Including F60–F69, F70–F79, F80–F89, F90–F99, Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4, and Z91.5 only.

(e) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

Table 2.20: Specialised care separations reported for Aboriginal and Torres Strait Islander patients,^(a) by RRMA category of area of usual residence, Australia, 1998-99

RRMA category of area of usual residence	Separations			Total patient days	Total psychiatric care days ^(c)	Average length of stay (overnight)	Psychiatric care days per overnight separation
	Same day	Overnight	Total				
Metropolitan	331	1,386	1,717	35,817	33,478	25.6	23.9
Rural	37	802	839	21,619	21,276	26.9	26.5
Remote	9	357	366	4,465	4,198	12.5	11.7
Not reported	6	133	139	3,515	3,353	26.4	25.2
Total	383	2,678	3,061	65,416	62,305	24.3	23.1

(a) A full definition of Aboriginal and Torres Strait Islander status is in the Glossary. The completeness of Aboriginal and Torres Strait Islander status data varies, hence these data should be used with caution.

(b) Rates are indirectly age-standardised based on the December 1998 Estimated Resident Population and the December 1998 Aboriginal and Torres Strait Islander population projections.

(c) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

Table 2.21: Separations with specialised psychiatric care by principal diagnosis in ICD-10-AM groupings^(a) and country of birth groups, Australia, 1998-99

Principal diagnosis	Australia		Other English-speaking		Non-English-speaking	
	Same day separations	Overnight separations	Separations per 1,000 population ^(b)		Separations per 1,000 population ^(b)	Separations per 1,000 population ^(b)
			separations	population ^(b)		
F00-F09 Organic, including symptomatic, mental disorders	98	916	0.1	1	217	0.1
F00-F03 Dementia	121	867	0.1	5	151	0.1
F04-F09 Other organic mental disorders	219	1,783	0.1	6	368	0.2
Total/						
F10-F19 Mental and behavioural disorders due to psychoactive substance use	2,197	3,127	0.4	358	472	0.5
F10 Mental behavioural disorders due to use of psychoactive substances	1,461	4,595	0.4	60	335	0.2
F11-F19 Mental behavioural disorders due to other psychoactive substances use	3,658	7,722	0.8	418	807	0.8
Total/						
F20-F29 Schizophrenia, schizotypal and delusional disorders	3,281	13,908	1.2	211	939	0.7
F20 Schizophrenia	1,970	6,490	0.6	458	583	0.6
F21-F29 Other schizophrenic, schizotypal, delusional disorders	5,251	20,398	1.8	669	1,522	1.3
Total/						
F30-F39 Mood (affective) disorders	22,685	22,282	3.1	2,954	2,890	3.6
F30-F39 Neurotic, stress-related and somatoform disorders	10,393	9,014	1.4	1,155	991	1.3
F40-F48 Behavioural syndromes associated with physiological disturbances and physical factors	4,322	1,188	0.4	55	92	0.1
F50-F59 Other mental health disorders ^(c)	4,429	5,298	0.7	318	446	0.5
Other principal diagnoses not related to mental health ^(a)	2,817	3,826	0.5	563	384	0.6
Not reported	16	620	0.0	3	56	0.0
Total/	53,790	72,131	8.8	6,141	7,556	8.4

(a) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition of a mental disorder outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.

(b) Rates are crude rates based on 31 December 1998 estimated resident population.

(c) Including F60-F69, F70-F79, F80-F89, F90-F99, Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5 only.

Overnight separations

separations

population

per 1,000

population

Table 2.22: Patient days and psychiatric care days^(a) of separations with specialised psychiatric care by country of birth groups, Australia, 1998–99



Principal diagnosis	Australia				Other English-speaking countries				Non-English-speaking countries			
	Patient days	Psychiatric care days per 1,000 ^(b)	Psychiatric care days per 1,000 ^(b)	Patient days	Psychiatric care days per 1,000 ^(b)	Psychiatric care days per 1,000 ^(b)	Patient days	Psychiatric care days per 1,000 ^(b)				
F00–F09 Organic, including symptomatic, mental disorders	63,368	54,404	3.79	15,034	9,779	6.00	30,594	24,090	8,80	9,651	3,53	
F00–F09 Dementia	80,328	76,943	5.36	4,605	2,783	1.71	10,799	9,651	8.80	33,741	12.33	
F04–F09 Other organic mental disorders	143,696	131,347	9.14	19,639	12,562	7.71	41,393	33,741	8.80	33,741	12.33	
<i>Total</i>												
F10–F19 Mental and behavioural disorders due to psychoactive substance use	101,182	83,669	5.82	5,868	4,978	3.05	19,653	19,033	6.96			
F10 Mental and behavioural disorders due to other psychoactive substances use	39,832	37,210	2.59	3,153	2,551	1.57	5,445	5,061	1.85			
<i>Total</i>	141,014	120,879	8.42	9,021	7,529	4.62	25,098	24,094	8.81			
F20–F29 Schizophrenia, schizotypal and delusional disorders	694,868	534,131	37.19	37,008	31,640	19.41	139,681	127,068	46.44			
F20 Other schizophrenic, schizotypal, delusional disorders	137,276	118,802	8.27	12,177	10,255	6.29	36,559	33,743	12.33			
<i>Total</i>	832,144	652,933	45.46	49,185	41,895	25.70	176,240	160,811	58.77			
F30–F39 Mood (affective) disorders	21,351	20,795	1.45	1,922	1,625	1.00	5,868	5,537	2.02			
F30 Manic episode	156,077	144,624	10.07	16,478	14,231	8.73	38,514	33,125	12.11			
F31 Bipolar affective disorders	256,534	231,402	16.11	33,353	26,770	16.42	79,421	70,767	25.86			
F32–F33 Depressive disorders	18,076	15,349	1.07	3,007	2,097	1.29	5,347	4,464	1.63			
F34–F39 Other mood (affective) disorders	452,038	412,170	28.69	54,760	44,723	27.44	129,150	113,893	41.62			
<i>Total</i>												
F40–F48 Neurotic, stress-related and somatoform disorders	94,838	80,343	5.59	10,184	7,158	4.39	19,190	17,061	6.23			
F50–F59 Behavioural syndromes associated with physiological disturbances and physical factors	30,069	27,377	1.91	1,629	1,378	0.85	3,130	2,892	1.06			
F50 Eating disorders												
F51–F59 Other behavioural syndromes associated with physiological disturbances, physical factors												
<i>Total</i>												
F60–F69 Disorders of adult personality and behaviour	4,119	4,003	0.28	490	471	0.29	914	907	0.33			
F70–F79 Mental retardation	34,188	31,380	2.18	2,119	1,849	1.13	4,044	3,799	1.39			
F80–F89 Disorders of psychological development	40,569	37,416	2.60	3,467	3,040	1.87	4,641	4,107	1.50			
F90–F98 Disorders onset usually occurring in childhood, adolescence	41,433	41,212	2.87	41	41	0.03	2,610	2,554	0.93			
F99 Mental disorder not otherwise specified	11,391	8,528	0.59	341	206	0.00	649	503	0.18			
Z03.2 Observation for suspected mental and behavioural disorder	829	827	0.06	29	29	0.02	36	36	0.01			
Other Z codes associated with mental disorders ^(c)	579	533	0.04	25	18	0.01	262	242	0.09			
Other principal diagnoses not related to mental health ^(d)	1,117	999	0.07	86	66	0.04	4,690	4,681	1.71			
Not reported	106,853	82,632	5.75	12,346	11,319	6.94	28,636	19,046	6.96			
<i>Total</i>	23,932	23,932	1.67	2,919	2,919	1.79	3,869	3,869	1.41			
	1,965,373	1,651,918	115.00	164,166	133,356	81.82	441,615	389,277	142.26			

(a) Does not include psychiatric care days from WA public acute or private hospitals. The total is therefore an underestimate of total psychiatric care days.

(b) Rates are crude rates based on 31 December 1998 estimated resident population data.

(c) Including Z09.3, Z13.3, Z13.4, Z54.3, Z76.5, Z91.4 and Z91.5 only.

(d) ICD-10-AM codes for each grouping are outlined in Appendix 1. Any disorders not falling within the definition outlined in Appendix 1 are included in the category 'Other principal diagnoses not related to mental health'.






Figure 2.6: Proportion of male and female separations in marital status groups, Australia, 1998–99, not including Victoria and private hospitals




Figure 2.7: Psychiatric care days per overnight separation by RRMA category of area of usual residence, public acute and private hospitals, Australia, 1998–99

Psychiatric care days




Figure 2.8: Psychiatric care days per overnight separation by RRMA category of area of usual residence, public psychiatric hospitals, Australia, 1998–99

Separations per 1,000 population




Figure 2.9: Separations with specialised psychiatric care by reported Aboriginal and Torres Strait Islander status and selected principal diagnosis in ICD-10-AM groupings per 1,000 population, Australia, 1998–99




Figure 2.10: Psychiatric care days per overnight separation with specialised psychiatric care, by reported Aboriginal and Torres Strait Islander status and selected principal diagnosis in ICD-10-AM groupings, Australia, 1998–99




Figure 2.11: Average number of psychiatric care days per overnight separation with specialised psychiatric care, by country of birth group and selected principal diagnosis in ICD-10-AM groupings, Australia, 1998–99