



Australian Government

Australian Institute of Health and Welfare



infocus

australian  
mesothelioma  
registry

# Mesothelioma in Australia 2022

Published November 2023

Australia has one of the highest measured incidence rates of mesothelioma in the world (Bray et al. 2017). Each year in Australia, between 700 and 800 people are diagnosed with the rare and aggressive cancer. In the 2019–20 financial year, the estimated health system expenditure for mesothelioma cases was \$32.1 million (total cancers \$12.1 billion) (AIHW 2022).

Men were more likely to be diagnosed with mesothelioma than women across all age groups, and the number of cases diagnosed each year for both men and women has steadily increased over the past 40 years. There is no cure for mesothelioma with the main cause being from exposure to asbestos – a material that has been banned in Australia since the end of 2003 (ASEA 2016). It can take many years after being exposed to asbestos (between 20 and 60 years) for mesothelioma to develop (Cancer Council 2019a).

This report presents the latest available statistics from the Australian Mesothelioma Registry (AMR), supplemented by data from the National Mortality Database (NMD) and the Australian Cancer Database (ACD).

Note that in this report, incidence data includes cases of mesothelioma notified up until 1 June 2023. This extraction date differs from previous reports. For further information see *Mesothelioma in Australia 2022 – methodology paper*.

## Main findings



**637 cases** of mesothelioma diagnosed in 2022 had been reported to the AMR as at 1 June 2023 – the **median age at diagnosis was 77**.



Between 1990–1994 and 2015–2019, the age-adjusted **relative survival of people with mesothelioma has increased**, most notably 1-year relative survival.



In 2021, **708 deaths of people** with mesothelioma (from any cause of death including mesothelioma) were recorded on the AMR – an age-standardised **mortality rate of 2.1 deaths per 100,000 population**.



More than **9 in 10** participants who completed an exposure assessment were assessed as having a history of **possible or probable exposure** to asbestos.

The AMR collects information on new cases of mesothelioma diagnosed in Australia since 1 July 2010. The Registry's main goals are to better understand the relationship between asbestos exposure and mesothelioma, to provide evidence in relation to current policies in how to deal with asbestos still in the environment, and to provide reliable information to policy makers and researchers. For more information on the AMR, see *Mesothelioma in Australia 2022 – methodology paper*.

AIHW

Stronger evidence,  
better decisions,  
improved health and welfare

## What is mesothelioma?

Mesothelioma is a form of cancer in the mesothelium – the protective lining on the inside of body cavities and the outside of internal organs. In 2022, 94% of cases reported to the AMR were pleural mesothelioma – that is, the cancer occurred in the mesothelial lining around the lungs. In the other 6% of cases, mesothelioma arose in other mesothelial-lined surfaces around the heart (pericardial), abdomen (peritoneal) and scrotal sac. More information on the diagnostic characteristics of cases of mesothelioma diagnosed in 2022 is available in Table A10 of [Mesothelioma in Australia 2022 – data tables](#). Although mesothelioma most commonly occurs in the chest, it is not a lung cancer and requires different forms of treatment (Cancer Council 2019a).

While mesothelioma is not curable, patients can undergo one or more different treatments to increase their survival period – including chemotherapy, radiotherapy, immunotherapy and surgery (Cancer Council 2019b). However, in the more advanced stages of the disease, the focus is on improving quality of life and easing symptoms (Cancer Council 2019c).

The main cause of mesothelioma is exposure to asbestos – a group of naturally occurring fibrous silicate materials with fibres invisible to the naked eye that can be inhaled into the lungs (enHealth 2013) where they do not readily break down.

### Australia's use of asbestos


Australia's consumption of asbestos peaked at a total of around 700,000 metric tonnes during the period 1970–1979 (Soeberg 2018; Leigh et al. 2002). Australia mined and imported asbestos, which was primarily used in the construction and transport industries because it was durable and resistant to fire and chemicals (ASEA 2016). Asbestos-related regulatory controls have been significantly tightened over time – asbestos-containing materials (ACMs) have been totally banned in Australia since December 2003, and it is now illegal to make, use or import them from another country. However, a large amount of asbestos still remains in older structures including houses, and products, potentially exposing workers and/or the public to asbestos if relevant safety procedures are not followed. The Commonwealth and state and territory work health and safety authorities have put in place strict asbestos-related regulatory measures to control asbestos exposure in workplaces (including safe management and removal of asbestos), to minimise asbestos-related diseases such as mesothelioma, lung cancer and asbestosis.

## How have the number of cases changed over time?

The AMR is the most up-to-date source of national data on mesothelioma incidence (number of new cases) in Australia, because case notifications are fast-tracked from state and territory cancer registries to the AMR. However, most notifications are still received in the year after diagnosis because of the time it takes to make a definitive diagnosis, and the time between diagnosis and notification to cancer registries and to the AMR. For more information about the challenges of collecting mesothelioma data and AMR processes, see page 17 of this report and [Mesothelioma in Australia 2022 – methodology paper](#).

Because not all cases of mesothelioma are reported to the AMR in the year that they are diagnosed, the number of cases recorded for each year continues to rise for all years dating back to 2011 (Figure 1). For example, of the 751 cases of mesothelioma diagnosed in 2015 to date and notified to the AMR, 67% of notifications were received in the year they were diagnosed (2015), 22% in 2016, and 11% were received from 2017 onwards.

**Figure 1: Difference in number of mesothelioma cases between originally published data<sup>1</sup> and current report during 2011 to 2021**




<sup>1</sup> Originally published data refers to the number of cases notified at the time those data were initially published.

Sources: AMR 2012, 2013, 2014, 2015, 2016 and 2017; AIHW 2018, 2019, 2020, 2021 and 2023a; AIHW analysis of AMR data as at 1 June 2023; Table A1 in *Mesothelioma in Australia 2022 – data tables*.

As at 1 June 2023, 637 cases of mesothelioma diagnosed in 2022 had been reported to the AMR (Figure 2). Men have been consistently more likely to be diagnosed with mesothelioma than women – this is expected because the majority of cases are from occupational exposure to asbestos in the type of environments in which men more commonly worked, such as the mining and construction industries. After adjusting for differences in the age structure of the populations, current available data show that the rate for persons averaged around 2.6 cases per 100,000 people between 2011 and 2022; for men, it averaged 4.6 per 100,000, and for women, it averaged 0.9 per 100,000.

**Figure 2: Number and age-standardised rate (ASR) (per 100,000 population) of people diagnosed with mesothelioma, by year and sex, 2011 to 2022**



Note: Rates have been age-standardised to the 2001 Australian Standard Population.


Source: AIHW analysis of AMR data as at 1 June 2023; Table A2 in *Mesothelioma in Australia 2022 – data tables*.

## Long-term trend

Reporting using AMR data is only possible for the period since July 2010 as that is when the data collection began. However, longer-term trends in the number and incidence rates of mesothelioma cases can be described by combining data from the ACD for 1982–2010 and the AMR for 2011–2022.

Between 1982 and 2022, the number of new cases of mesothelioma reported annually steadily increased – from 135 to 504 for men and from 22 to 133 for women (Figure 3). To date, the year in which the highest overall number of cases (826) were diagnosed is 2017. It is important to note that the apparent fall in cases in 2022 is likely due to delays in the AMR receiving notifications – the number of cases for 2022 are expected to rise after 1 June 2023 (date these data were extracted). Also note that the number of cases may continue to increase slightly for all years, and particularly from 2017 onwards, as more cases are notified to the AMR (see Table A1 in *Mesothelioma in Australia 2022- data tables* for further information).

Figure 3: Number of people diagnosed with mesothelioma, by year and sex, 1982 to 2022




Note: Data for 1982–2010 sourced from ACD; Data for 2011–2022 sourced from AMR.

Sources: 1982–2010: AIHW 2023b; 2011–2022: AIHW analysis of AMR data as at 1 June 2023; Table A3 in *Mesothelioma in Australia 2022 - data tables*.

Between 1982 and 2017, the crude incidence rate rose for men from 1.8 to 5.6 cases per 100,000 population. For women, the rate rose from 0.3 in 1982 to 1.4 cases per 100,000 population in 2018 (Figure 4). The crude incidence rate for persons rose from 1.0 in 1982 to a peak of 3.4 cases per 100,000 population in 2017 and has since remained at around 3.1 cases per 100,000 population.

**Figure 4: Crude rate (per 100,000 population) of people diagnosed with mesothelioma, by year and sex, 1982 to 2022**




Note: Data for 1982–2010 sourced from ACD; Data for 2011–2022 sourced from AMR.

Sources: 1982–2010: AIHW 2023b; 2011–2022: AIHW analysis of AMR data as at 1 June 2023; Table A3 in *Mesothelioma in Australia 2022 – data tables*.

After adjusting for differences in the age structure of the populations, the age-standardised incidence rate rose between 1982 and 2003 for both men (2.3 to 5.9 cases per 100,000 population) and women (0.3 to 1.0 cases per 100,000 population) (Figure 5). The incidence rate for persons rose from 1.2 in 1982 to a peak of 3.2 cases per 100,000 population in 2003 and has since remained at around 2.8 cases per 100,000 population.

**Figure 5: Age-standardised rate (per 100,000 population) of people diagnosed with mesothelioma, by year and sex, 1982 to 2022**



Notes:

- Rates have been age-standardised to the 2001 Australian Standard Population.
- Data for 1982–2010 sourced from ACD; Data for 2011–2022 sourced from AMR.


Sources: 1982–2010: AIHW 2023b; 2011–2022: AIHW analysis of AMR data as at 1 June 2023; Table A3 in *Mesothelioma in Australia 2022 – data tables*.

## How does mesothelioma diagnosis vary by age and sex?

The age of people in the AMR who were diagnosed in 2022 ranged from 43 to 98. The median age at diagnosis was 77, and a greater number of men than women were diagnosed across all age groups. Figure 6 shows the number of people diagnosed with mesothelioma by age group and sex.

Mesothelioma has a long and highly varied latency period (Marinaccio et al. 2007; Shavelle et al. 2017), with symptoms typically appearing decades after a person has been exposed to asbestos. An Australian study by Reid et al. (2014) that analysed the data of more than 22,000 people who had been exposed to asbestos found that for those with mesothelioma, the median time since first exposure to diagnosis was 38 years. A similar study by Olsen et al. (2011) based on data from people diagnosed in Western Australia between 1960 and 2008 reported a latency period of between 33 and 44 years. Most mesothelioma diagnoses are made later in life.


Figure 6: Number of people diagnosed with mesothelioma, by age group and sex, 2022



Source: AIHW analysis of AMR data as at 1 June 2023; Table A4 in *Mesothelioma in Australia 2022 – data tables*.

In 2022, age-specific mesothelioma incidence rates (the number of new mesothelioma cases per 100,000 population in specific age groups) generally increased with increasing age – for age 85 and over, the rates for men were 41.4 cases per 100,000 population and for women were 6.0 cases per 100,000 population. Men consistently had higher rates than women across all age groups (Figure 7).

**Figure 7: Rate (per 100,000 population) of people diagnosed with mesothelioma, by age group and sex, 2022**



Source: AIHW analysis of AMR data as at 1 June 2023; Table A4 in *Mesothelioma in Australia 2022 – data tables*.

## How do rates vary by state and territory?

Because of the small number of people diagnosed in many age groups in some states and territories, data have been grouped for the years 2019–2022. This enables the data to be directly age standardised to remove the effect of differing age structures between the populations, which can affect incidence rates. During the period 2019–2022, age-standardised rates of mesothelioma ranged from 0.4 cases per 100,000 people in the Northern Territory to 3.0 cases per 100,000 people in Western Australia (Table 1) – likely due to extensive asbestos mining in the past (Cancer Council 2019a). Additional data are available for 2015–2018 in Table 1 in *Mesothelioma in Australia 2022 – data tables*.

**Table 1: Number and age-standardised rate (per 100,000 population) of people diagnosed with mesothelioma, by sex and state/territory, 2019–2022**

State of diagnosis	Males		Females		Persons	
	No.	Rate	No.	Rate	No.	Rate
NSW	754	3.8	176	0.8	930	2.2
Vic	517	3.4	141	0.8	658	2.0
Qld	536	4.4	137	1.0	673	2.6
WA	311	5.1	81	1.2	392	3.0
SA	186	3.8	50	0.9	236	2.2
Tas	65	4.0	12	0.7	77	2.2
ACT	22	2.8	11	1.1	33	1.8
NT	2	0.4	2	0.4	4	0.4
Australia	2,393	3.9	610	0.9	3,003	2.3

Note: Due to small counts in some states and territories, data have been grouped into the years 2019–2022, to enable rates to be directly age-standardised to the 2001 Australian Standard Population.

Source: AIHW analysis of AMR data as at 1 June 2023.

# How long do people live after diagnosis?

## What is relative survival?

Relative survival is a measure of the survival of people with mesothelioma compared with that of the general population. It is calculated by dividing the observed survival of people with mesothelioma by expected survival of the general population, where the numerator and denominator have been matched for age, sex and calendar year.


The population diagnosed with mesothelioma continues to change over time and is more commonly older in more recent years, averaging around 75 years of age at diagnosis between 2011 and 2022 (AMR data). Even though relative survival for people with mesothelioma is improving over time, the extent of improvement reflected in relative survival rates is diminished because an increasing proportion of people with mesothelioma are older in more recent years than in the past and survival rates for older people tend to be lower. An analysis of mesothelioma using ACD data and AMR data shows that median age at diagnosis has increased from 63 years in 1982 to 77 years in 2022 (AIHW 2023b). By age-adjusting to a group with a consistent age structure, such as those people with mesothelioma in 2015–2019, age-adjusted relative survival rates account for the impact different age structure can have on comparisons over time in relative survival rates.

For further information see [Mesothelioma in Australia 2022 – methodology paper](#).

Due to its aggressive nature, mesothelioma has a very low survival rate compared to other cancers (AIHW 2023b). The condition is often diagnosed at advanced stages, because early symptoms can go unnoticed or be mistaken as symptoms for other conditions or diseases (ADRI 2019; Cancer Council 2019a). Many factors can affect the person's chances of survival, including their age at diagnosis, their overall health status and the type of mesothelioma they have.

Linked data from the ACD and National Death Index (NDI), shows that the average length of time that people survive after being diagnosed with mesothelioma is gradually improving over time. Figure 8 shows the 1, 3 and 5-year age-adjusted relative survival of people with mesothelioma from 1990–1994 to 2015–2019 – all survival rates have increased over this period, most notably the 1-year survival. This apparent improvement in survival may be due to factors such as diagnosing mesothelioma at an earlier stage and improvements over time in how patients are treated.

**Figure 8: 1, 3 and 5-year age-adjusted relative survival of people diagnosed with mesothelioma, 1990–1994 to 2015–2019**



**Notes**


1. Relative survival was calculated using the period method. For more information on survival calculations, see the "Using the data" and "Technical Notes" pages within the Cancer Data in Australia web report.
2. Records with unknown age at diagnosis and/or those who identify as intersex, non-binary or other were excluded from the analysis.
3. Age-adjusted backward-looking: The age composition of those diagnosed with the selected cancer group/site in earlier periods is adjusted to equal the age composition of those diagnosed with the selected cancer in the 2015–2019 survival period.
4. Please read [Cancer data commentary number 6](#) for more information about age-adjusted survival rates.

Source: AIHW 2023b; Table A5 in *Mesothelioma in Australia 2022 – data tables*.

## Death rate

Using the NMD to report on long-term mortality trends, the number of mesothelioma deaths has increased from 416 in 1997 to 708 in 2021. Deaths from mesothelioma increased generally but with fluctuations over the period among both men and women. For men, the number of deaths in 2021 was 1.6 times higher than in 1997, while for women there were over twice as many deaths in 2021 compared to 1997 (Figure 9).

Figure 9: Number of deaths from mesothelioma, by year and sex, 1997 to 2021



### Notes


- Figures are presented by year of death for 1997–2020 and year of registration of death for 2021.
- Deaths registered in 2019 and earlier are based on the final version of Cause of Death Unit Record Files (CODURF). Deaths registered in 2020 are based on the revised version and deaths registered in 2021 are based on the preliminary version. Revised and preliminary versions are subject to further revision by the Australian Bureau of Statistics (ABS). For further information see [Mesothelioma in Australia 2022 – methodology paper](#).
- Rates have been age-standardised to the 2001 Australian Standard Population.

Source: AIHW 2023b; Table A6 in [Mesothelioma in Australia 2022 – data tables](#).

Although the number of deaths has increased over time, the age standardised rate of deaths has remained fairly stable (Figure 10). The most likely driver for the increasing number of deaths is Australia's increasing and ageing population (whereby older Australians are accounting for an increasing proportion of the population) (ABS 2019).

From 1997 to 2021, age-standardised mesothelioma mortality rates fluctuated between 2.1 and 2.7 deaths per 100,000 population (Figure 10). Rates for men ranged from 3.6 deaths per 100,000 in 2021 to 5.0 deaths per 100,000 population in 2001 and 2004. Rates for women fluctuated around 0.8 deaths per 100,000 over the period.

**Figure 10: Age-standardised rate (per 100,000 population) of deaths from mesothelioma, by year and sex, 1997 to 2021**



#### Notes

- Figures are presented by year of death for 1997–2020 and year of registration of death for 2021.
- Deaths registered in 2019 and earlier are based on the final version of Cause of Death Unit Record Files (CODURF). Deaths registered in 2020 are based on the revised version and deaths registered in 2021 are based on the preliminary version. Revised and preliminary versions are subject to further revision by the Australian Bureau of Statistics (ABS). For further information see [Mesothelioma in Australia 2022 – methodology paper](#).
- Rates have been age-standardised to the 2001 Australian Standard Population.

Source: AIHW 2023b; Table A6 in [Mesothelioma in Australia 2022 – data tables](#).

## Asbestos exposure among people with mesothelioma

Previous research has shown mesothelioma can be associated with occupational and non occupational exposure to asbestos. Historically, occupational exposure has been dominated by asbestos mining, manufacturing and use of asbestos-containing materials (enHealth 2013; Safe Work Australia 2014). Because mesothelioma typically develops a long time after exposure, the majority of current cases in this report are probably related to occupational exposure in workplaces that occurred before current occupational asbestos regulations and practices came into effect. The results here should not be interpreted as indicative of current risk in workplaces today.

In total, 1,413 people (1,134 men and 279 women) diagnosed with mesothelioma since 1 July 2010 consented to participate in the voluntary asbestos **exposure assessment** as at 1 June 2023. Of these, 1,218 people (966 men and 252 women) completed both the questionnaire and telephone interview components of the assessment.

Based on the jobs held by the participant during their working life, relevant job-specific questionnaire modules were allocated for the participant's telephone interview. For example, participants who have worked in jobs such as electrician, plumber and carpenter may be allocated the 'Trades' module. Participants could be allocated multiple job-specific modules for different jobs. Participants' lifetime exposure in non-occupational settings (such as their home) was also evaluated with a non-occupational module. For the purposes of this assessment, potential exposures to asbestos were then classified according to the likelihood that they were above background levels of 0.0001 f/ml (fibres of asbestos per millilitre) (Brown 2001). **Probability of exposure** was assessed as either '**probable**', '**possible**' or '**unlikely**', and **level of exposure** as either '**high**', '**medium**' or '**low**'. For more information, see [Mesothelioma in Australia 2022 – methodology paper](#).

Participants' lifetime exposure in non-occupational settings (such as their home) was also evaluated with a non-occupational module. For the purposes of this assessment, potential exposures to asbestos were then classified according to the likelihood that they were above background levels of 0.0001 f/ml (fibres of asbestos per millilitre) (Brown 2001). **Probability of exposure** was assessed as either '**probable**', '**possible**' or '**unlikely**', and **level of exposure** as either '**high**', '**medium**' or '**low**'. For more information, see [Mesothelioma in Australia 2022 – methodology paper](#).

## Nine in ten participants were assessed as having some exposure to asbestos

More than 9 in 10 (94% or 1,141) people were assessed as having possible or probable exposure to asbestos (Table 2). For men, this exposure most commonly occurred in their jobs. Occupational exposure to asbestos is typically associated with greater exposure and frequency than non-occupational exposure.

Of the 1,141 people for whom exposure was detected, most men had some form of occupational exposure, whereas very few women did:

- 78% (716) of men provided information indicating possible or probable occupational exposure ('occupational exposure only' and 'both occupational and non-occupational exposure'), compared with 6.8% (15) of women (Table 2).
- 99% (211) of women provided information indicating non-occupational exposure ('non occupational exposure only' and 'both occupational and non-occupational exposure'), compared with 86% (789) of men (Table 2).

**Table 2: Occupational and non-occupational exposure assessment, by sex, 2010–2022**

Any exposure indicated	Men		Women		Persons	
	No.	%	No.	%	No.	%
Occupational exposure only	130	14.1	1	0.5	131	11.5
Non-occupational exposure only	203	22.1	207	93.2	410	35.9
Both occupational and non-occupational exposure	586	63.8	14	6.3	600	52.6
<b>Total</b>	<b>919</b>	<b>100</b>	<b>222</b>	<b>100</b>	<b>1,141</b>	<b>100</b>

Note: Of the 1,218 participants, 77 (6.3%) participants (47 men and 30 women) were assessed as having neither occupational nor non-occupational exposure. Although it was not possible to identify asbestos exposure among these participants, this should not be taken to mean that these participants have never been exposed to asbestos; rather, it means that no evidence of exposure above background was obtained by the exposure assessment methods used.

Source: AIHW analysis of AMR data as at 1 June 2023, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2022.

Table 3 shows the results from the asbestos exposure assessment among the 5 states and territories with the largest number of participants; New South Wales, Victoria, Queensland, Western Australia and South Australia. There were insufficient numbers of participants to present results for Northern Territory, Australian Capital Territory and Tasmania, so these were excluded.

Table 3 shows that the proportion of participants assessed as having:

- 'occupational exposure only' ranged from 11% to 13% across states
- 'non-occupational exposure only' ranged from 29% to 43% across states
- 'both occupational and non-occupational exposure' ranged from 45% to 60% across states.

As these percentages are based on small numbers of participants, drawing conclusions about apparent differences between states is not recommended.

**Table 3: Occupational and non-occupational exposure assessment, by state, 2010–2022**

Any exposure indicated	State									
	NSW		Vic		Qld		WA		SA	
	No.	%	No.	%	No.	%	No.	%	No.	%
Occupational exposure only	49	12.0	27	11.9	20	10.6	19	10.1	11	13.1
Non-occupational exposure only	145	35.6	98	43.4	55	29.3	61	32.4	35	41.7
Both occupational and non-occupational exposure	213	52.3	101	44.7	113	60.1	108	57.4	38	45.2
<b>Total</b>	<b>407</b>	<b>100</b>	<b>226</b>	<b>100</b>	<b>188</b>	<b>100</b>	<b>188</b>	<b>100</b>	<b>84</b>	<b>100</b>

#### Notes

1. Of the 1,218 participants, 77 (6.3%) participants (47 men and 30 women) were assessed as having neither occupational nor non-occupational asbestos exposure. Although it was not possible to identify asbestos exposure among these participants, this should not be taken to mean that these participants have never been exposed to asbestos; rather it means that no evidence of exposure above background was obtained by the exposure assessment methods used.
2. As these percentages are based on small numbers of participants, drawing conclusions about apparent differences between states is not recommended.
3. There were insufficient numbers of participants to present results for NT, ACT and Tasmania.


Source: AIHW analysis of AMR data as at 1 June 2023, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2022.

## Occupational asbestos exposure

Figure 11 shows the estimated likelihood of exposure among jobs assessed using the three most commonly allocated job-specific interview modules. Based on the interview data collected, 81% of participants who received the 'Trades' module, 74% of those who received the 'Water transport' module, and 38% of participants who received the 'Land transport' module were assessed as having had 'probable' exposure to asbestos for jobs in those categories.

Results are based on the responses people gave to each specific module - participants are typically assigned multiple job-specific modules for different jobs, so the sum of respondents assigned to each module does not equal the total number of participants. Because questionnaire modules are assigned only to jobs with some likelihood of exposure, the finding of exposure in a high proportion of the jobs that respondents were questioned about is to be expected.

**Figure 11: Exposure assessment results by most commonly used job-specific modules, 2010–2022**



Note: Bracketed figures (p) are the number of participants allocated this job-specific module at least once.

Source: AIHW analysis of AMR data as at 1 June 2023, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2022; Table A7 in *Mesothelioma in Australia 2022 – data tables*.

## Occupational exposure by job type

For participants who received the same module for more than one job, the results presented are based on a participant's highest exposed job in that job category. Because participants can have different exposure probabilities and/or levels for different jobs, the exposure estimate reported here is the maximum exposure likely in that job category for each individual. For example, if a participant had two trade-related jobs which were assessed as having a different probability and/or level of exposure, the higher of the two would be reported for that participant in that job category. Participants may be assigned different modules for different types of jobs, so the sum of respondents assigned to each module does not equal the total number of participants.


Jobs are coded according to the Australian and New Zealand Standard Classification of Occupations (ANZSCO 2013, Version 1.3).

Among participants who received the 'Trades' module, 'possible or probable' exposure to asbestos was assessed in:

- 94% of the 93 people in electrical trade jobs
- 90% of the 267 people in building trade jobs
- 90% of the 60 people in metal production and fabrication jobs
- 86% of the 56 people in plumber/gasfitter jobs
- 79% of the 101 people in metal fitter/turner and toolmaker jobs

In contrast, only 50% of the 18 people in telecommunication technician jobs and 70% of the 20 people in engineer jobs were assessed as 'possible or probable' exposure to asbestos (see Figure 12).

**Figure 12: Occupational asbestos exposure by job title for the 'Trades' module, 2010–2022**



Note: Bracketed figures (p) are the number of participants allocated this job title at least once.


Source: AIHW analysis of AMR data as at 1 June 2023, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2022; Table A8 in *Mesothelioma in Australia 2022 – data tables*.

Among participants who received the 'Land or Water transport' modules, 'possible or probable' exposure to asbestos was assessed in:

- All of the 32 people in marine engineer jobs
- 90% of the 29 people in shipwright/boatbuilder jobs
- 80% of the 5 people in stevedore/waterside worker jobs
- 71% of the 59 people in automotive maintenance jobs

In contrast, only 20% of the 66 people in other land transport jobs were assessed as 'possible or probable' exposure to asbestos (see Figure 13). For more information, see *Mesothelioma in Australia 2022 – methodology paper*.

**Figure 13: Occupational asbestos exposure by job title for the ‘Water transport’ and ‘Land transport’ modules, 2010–2022**



Note: Bracketed figures (p) are the number of participants allocated this job title at least once.

Source: AIHW analysis of AMR data as at 1 June 2023, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2022; Table A8 in *Mesothelioma in Australia 2022 – data tables*.

## Non-occupational asbestos exposure

All 1,218 participants completed the non-occupational questionnaire module. Of those where exposure was indicated, 1,010 (89%) were assessed as having had ‘possible or probable’ exposure in non-occupational contexts. Non-occupational exposure to asbestos is typically associated with lower exposure and frequency than occupational exposure. It was common to have indications of exposure in more than one non-occupational context, so a number of participants are counted in more than one category. Therefore, the sum of participants in each category does not add up to the total number of participants. The most common contexts in which non-occupational asbestos exposure was assessed as possible or probable were among those who reported ever having:

- undertaken major home renovations that involved asbestos products (excluding paid work) (51% assessed as possible or probable exposure)
- lived in a house undergoing renovations (39% assessed as possible or probable exposure)
- serviced car brakes/clutch (excluding paid work) (30% assessed as possible or probable exposure)
- lived in the same home as someone with a job where they were exposed to asbestos and who came home dusty (20% assessed as possible or probable exposure).
- lived in a house made of fibro that was built between 1947 and 1987 (12% assessed as possible or probable exposure).

See Table A9 in *Mesothelioma in Australia 2022 – data tables* for further information on sources of non-occupational exposure.

## No asbestos exposure

For 77 participants (6.3% of all 1,218 participants – 47 men and 30 women), there was no indication of asbestos exposure above background levels in either occupational or non-occupational contexts. This does not mean that these participants have never been exposed to asbestos; rather it means that no evidence of exposure above background levels was obtained using the exposure assessment methods.

### What are the challenges in collecting data and reporting on mesothelioma and associated asbestos exposure?

Confirming a diagnosis of mesothelioma is often very challenging for a variety of reasons:

- Symptoms of mesothelioma are common to many other conditions, and mesothelioma cells can often look similar to cells of other cancers (Cancer Council 2019b).
- Because mesothelioma symptoms are not specific to the condition, diagnosis is often complicated. Diagnostic confirmation of mesothelioma generally involves a number of clinical investigations, including biopsies, radiology and clinical examinations conducted by a multidisciplinary team (van Zandwijk et al. 2013).
- Diagnostic and treatment practices for mesothelioma are not equally distributed across Australia (van Zandwijk et al. 2013).

If a mesothelioma diagnosis is uncertain for any reason, the AMR is not notified and the case remains unrecorded, until such a time as the diagnosis is confirmed. Other challenges in collecting data and reporting on mesothelioma include:

- Although state and territory cancer registries fast-track mesothelioma notifications, there is still a time lag between a person's diagnosis, their inclusion in the AMR data set and (if consent is given) when they are interviewed for the AMR's asbestos exposure collection. Reasons for this lag include the time it takes to make a definitive diagnosis and the time between diagnosis and notification to cancer registries and to the AMR. Case verification and recruitment processes also vary between state and territory cancer registries.
- The participation rate in the asbestos exposure assessment of the AMR is low, at around 15% of people with mesothelioma. This is partially due to people dying or being too unwell to participate.

## Where do I go for more information?

More information on the AMR is available at <https://mesothelioma-australia.com/home>. The report *Mesothelioma in Australia 2022* and previous annual reports are available at <https://www.mesothelioma-australia.com/publications-and-data/publications>. People diagnosed with mesothelioma can choose to self-notify by contacting the AMR via email at amr@aihw.gov.au or via the toll-free information line on 1800 378 861.

For more information about the regulation and eradication of asbestos in Australia visit the Safe Work Australia website at <https://www.swa.gov.au> and the Asbestos Safety and Eradication Agency website Asbestos Safety and Eradication Agency: <https://www.asbestossafety.gov.au/>.

# Glossary

**age-adjusted survival:** A method to remove the influence of changes in the ages of those diagnosed with a specific cancer type (or group) over time when considering changes in relative survival rates over time. The population used in this report is people diagnosed with mesothelioma in the latest period, which was used to adjust the survival rates for all the previous years.

**age-standardisation:** A way to remove the influence of age when comparing populations with different age structures. This is usually necessary because the rates of many diseases vary strongly with age (usually increasing with increasing age). The age structures of the different populations are converted to the same 'standard' structure and then the disease rates that would have occurred with that structure are calculated and compared. This report uses the direct method of age standardisation.

**age-standardised rate:** A rate that results from removing the influence of age by converting the age structures of the different populations to the same 'standard' structure, providing a more valid way of comparing rates from populations with different age structures. The population used in this report is the 2001 standard population.

**age-specific rate:** The rate for a specific age-group. The numerator and denominator pertain to the same age group.

**Australian and New Zealand Standard Classification of Occupations (ANZSCO):** is primarily a statistical classification designed to aggregate and organise data collected about jobs or individuals and is based on the skill level and specialisation usually necessary to perform the tasks of the specific occupation, or of most occupations in the group.

**Australian Cancer Database (ACD):** Contains data on all new cases of cancer diagnosed in Australia (except basal and squamous cell carcinomas of the skin) since 1982.

**Cause of death unit record file (CODURF):** Is a dataset containing information relating to all deaths registered in Australia for a given reference year.

**crude rate:** the number of events in a given period divided by the size of the population at risk in a specified time period.

**exposure assessment:** a voluntary component of the AMR whereby patients provide their occupational and residential histories and are interviewed using a structured modular questionnaire. Information collected from participating patients is assessed via the Occupational Integrated Database Exposure Assessment System (OccIDEAS) which provides an indication of possible asbestos exposure.

**incidence:** The number of new cases (of an illness or event, and so on) occurring during a given period, often expressed as a rate (number per population).

**mortality:** The number or rate of deaths in a population during a given time period.

**National Death Index (NDI):** A catalogue of death records used in data linkage for epidemiological studies.

**National Mortality Database (NMD):** Holds records for all deaths in Australia since 1964.

**non-occupational exposures:** Chemical, biological, psychosocial, physical and other factors from places other than the workplace that can potentially cause harm. Examples include contact with asbestos during private house renovations and living in the same home as someone with an asbestos-exposed occupation who came home dusty.

**occupational exposures:** Chemical, biological, psychosocial, physical and other factors in the workplace that can potentially cause harm.

**relative survival:** the probability of being alive for a given amount of time after diagnosis compared with the general population. A 5-year relative survival figure of 100% means that the cancer has no impact on the person's chance of still being alive 5 years after diagnosis, whereas a figure of 50% means that the cancer has halved that chance.

**underlying cause of death:** The disease or injury that initiated the sequence of events leading directly to death.

## References

- ABS (Australian Bureau of Statistics) (2019) *Australian Historical Population Statistics, 2016*, ABS, Australian Government, accessed 19 July 2022.
- AIHW (Australian Institute of Health and Welfare) (2018) *Mesothelioma in Australia 2017*, AIHW, Australian Government, accessed 19 July 2022.
- AIHW (2019) *Mesothelioma in Australia 2018*, AIHW, Australian Government, accessed 19 July 2022.
- AIHW (2020) *Mesothelioma in Australia 2019*, AIHW, Australian Government, accessed 19 July 2022.
- AIHW (2021) *Mesothelioma in Australia 2020*, AIHW, Australian Government, accessed 19 July 2022.
- AIHW (2022) *Disease expenditure in Australia 2019–20*, AIHW, Australian Government, accessed 16 January 2023.
- AIHW (2023a) *Mesothelioma in Australia 2021*, AIHW, Australian Government, accessed 11 July 2023.
- AIHW (2023b) *Cancer data in Australia*, AIHW, Australian Government, accessed 31 August 2023.
- AMR (Australian Mesothelioma Registry) (2012) *Australian Mesothelioma Registry 1st annual report: Mesothelioma in Australia 2011*. Cancer Institute NSW, NSW Health, accessed 19 July 2022.
- AMR (2013) *Australian Mesothelioma Registry 2nd annual report: Mesothelioma in Australia 2012*, Cancer Institute NSW, NSW Health, accessed 19 July 2022.
- AMR (2014) *Australian Mesothelioma Registry 3rd annual report: Mesothelioma in Australia 2013*, Cancer Institute NSW, NSW Health, accessed 19 July 2022.
- AMR (2015) *Australian Mesothelioma Registry 4th annual report: Mesothelioma in Australia 2014*, Cancer Institute NSW, NSW Health, accessed 19 July 2022.
- AMR (2016) *Australian Mesothelioma Registry 5th annual report: Mesothelioma in Australia 2015*, Cancer Institute NSW, NSW Health, accessed 19 July 2022.
- AMR (2017) *Australian Mesothelioma Registry 6th annual report: Mesothelioma in Australia 2016*, Cancer Institute NSW, NSW Health, accessed 19 July 2022.
- ADRI (Asbestos Diseases Research Institute) (2019) *Mesothelioma*, Asbestos Diseases Research Institute, accessed 19 July 2022.
- ASEA (Asbestos Safety and Eradication Agency) (2016) *Asbestos Safety and Eradication Agency Reports 2016: Asbestos: the International and Australian Contexts*, ASEA, Australian Government, accessed 12 July 2022.
- Bray F, Colombet M, Mery L, Piñeros M, Znaor A, Zanetti R and Ferlay J (2017) 'Cancer incidence in five continents Vol. XI', *International Agency for Research on Cancer*, Viewed 4 July 2021, <http://ci5.iarc.fr>.
- Brown SK (2001) 'Asbestos', In: Spengler JD, Samet JMS and McCarthy JF (eds) *Indoor air quality handbook*, McGraw-Hill Education, 1-17.
- Burgers JA and Damhuis RA (2004) 'Prognostic factors in malignant mesothelioma', *Lung Cancer*, 45 Supplement 1:S49–S54, <https://doi.org/10.1016/j.lungcan.2004.04.012>.
- Cancer Council (2019a) *Mesothelioma*, Cancer Council, accessed 19 July 2022.
- Cancer Council (2019b) *Trimodality therapy for pleural mesothelioma*, Cancer Council, accessed 19 July 2022.
- Cancer Council (2019c) *Diagnosing mesothelioma*, Cancer Council, accessed 19 July 2022.
- enHealth (Environmental Health Standing Committee) (2013) Asbestos: a guide for householders and the general public, Australian Health Protection Principal Committee, accessed 19 July 2022.
- Leigh J, Davidson P, Hendrie L and Berry D (2002) 'Malignant mesothelioma in Australia, 1945–2000', *American Journal of Industrial Medicine*, 41:188–201, <https://doi.org/10.1002/ajim.10047>.
- Marinaccio A, Binazzi A, Cauzillo G, Cavone D, De Zotti R, Ferrante P, Gennaro V, Gorini G, Menegozzo M, Mensi C and Merler E (2007) 'Analysis of latency time and its determinants in asbestos related malignant mesothelioma cases of the Italian register', *European Journal of Cancer*, 43(18):2722–2728, <https://doi.org/10.1016/j.ejca.2007.09.018>.
- Olsen NJ, Franklin PF, Reid, A, de Klerk NH, Threlfall TJ, Shilkin K and Musk B (2011) 'Increasing incidence of Mesothelioma in Australia 2022

malignant mesothelioma after exposure to asbestos during home maintenance and renovation', *The Medical Journal of Australia*, 195(5): 271–274, <https://doi.org/10.5694/mja11.10125>.

Reid A, de Klerk NH, Magnani C, Ferrante D, Berry G, Musk AW and Merler E (2014) 'Mesothelioma risk after 40 years since first exposure to asbestos: a pooled analysis', *Thorax*, 69:843–850, <http://dx.doi.org/10.1136/thoraxjnl-2013-204161>.

Safe Work Australia (2014) *Asbestos-related disease indicators*. Safe Work Australia, Australian Government, accessed 19 July 2022.

Shavelle R, Vavra-Musser K, Lee K and Brookes J (2017) 'Life Expectancy in Pleural and Peritoneal Mesothelioma', *Lung Cancer International*, doi: 10.1155/2017/2782590.

Soeberg MJ, Vallance DA, Keena V, Takahashi K and Leigh J (2018) 'Australia's ongoing legacy of asbestos: significant challenges remain even after the complete banning of asbestos almost fifteen years ago', *International Journal of Environmental Research and Public Health*, 15:384. doi:10.3390/ijerph15020384.

van Zandwijk N, Clarke C, Henderson D, Musk AW, Fong K, Nowak A, Loneragan R, McCaughan B, Boyer M, Feigen M and Currow D (2013) 'Guidelines for the diagnosis and treatment of malignant pleural mesothelioma', *Journal of Thoracic Disease*, 5(6): E254–E307, <https://doi.org/10.3978%2Fj.issn.2072-1439.2013.11.28>.

## Acknowledgments

This report was prepared by Mr Tom Watson, Ms Sue Barker and Mr Brett Davis from the AMR team at the AIHW. The AMR team would also like to acknowledge the valuable guidance and advice from Mr Justin Harvey and Mr Richard Juckes. The AMR Expert Advisory Group also provided valuable advice: Professor Tim Driscoll (Chair), Associate Professor Fraser Brims, Ms Elizabeth Chalker, Dr Claire Cooke Yarborough, Ms Justine Ross, Professor Karen Walker-Bone and Associate Professor Sonja Klebe.

The AIHW acknowledges the data and assistance provided by all state and territory cancer registries, and thanks staff at the Monash Centre for Occupational and Environmental Health for undertaking and reporting on the asbestos exposure component of the AMR, and staff at the Social Research Centre for their work interviewing people diagnosed with mesothelioma.

The AMR is funded by Safe Work Australia.

---

© Australian Institute of Health and Welfare 2023 

All material presented in this document is provided under a Creative Commons Attribution 4.0 International licence, with the exception of the Commonwealth Coat of Arms (the terms of use for the Coat of Arms are available at <https://www.pmc.gov.au/government/commonwealth-coat-arms>) or any material owned by third parties, including for example, design, layout or images obtained under licence from third parties and signatures. All reasonable efforts have been made to identify and label material owned by third parties.

The details of the relevant licence conditions are available on the Creative Commons website (available at <https://creativecommons.org>), as is the full legal code for the CC BY 4.0 license.

### Suggested citation

Australian Institute of Health and Welfare (2023) *Mesothelioma in Australia 2022*, catalogue number CAN 156, AIHW, Australian Government.

ISBN 978-1-923085-16-26 (Online)  
ISBN 978-1-923085-16-27 (Print)  
DOI: 10.25816/mhy7-bw34

Any enquiries about copyright should be directed to the Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601, Tel: (02) 6244 1000, Email: [info@aihw.gov.au](mailto:info@aihw.gov.au).