

Better information and statistics for better health and wellbeing

Monitoring the impact of air pollution on asthma in Australia

A methods paper

February 2010

Australian Institute of Health and Welfare Canberra

Cat. no. ACM 18

The Australian Institute of Health and Welfare is Australia's national health and welfare statistics and information agency. The Institute's mission is better information and statistics for better health and wellbeing.

© Australian Institute of Health and Welfare 2010

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Australian Institute of Health and Welfare. Requests and enquiries concerning reproduction and rights should be directed to the Head, Media and Communications Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.

A complete list of the Institute's publications is available from the Institute's website <www.aihw.gov.au>.

ISBN 978-1-74249-007-6

Suggested citation

Australia: a methods paper. Cat. no. ACM 18. Canberra: AIHW.

Australian Institute of Health and Welfare

Board Chair

Hon. Peter Collins, AM, QC

Director

Penny Allbon

Any enquiries about or comments on this publication should be directed to:

Dr Adrian Webster

Australian Institute of Health and Welfare

GPO Box 570

Canberra ACT 2601

Phone: (02) 6244 1119

Email: adrian.webster@aihw.gov.au

Published by the Australian Institute of Health and Welfare

Please note that there is the potential for minor revisions of data in this report. Please check the online version at <www.aihw.gov.au> for any amendments.

Contents

Ac	knowledgments	v
Su	mmary	vi
1	Introduction	1
	Components of air pollution	2
	Health effect estimates	3
	Air pollution and incidence data	5
2	Key challenges	7
	Regional variation	7
	Time and lag effects	7
	Measurement methods	8
	Age ranges	8
	Baseline incidence	8
	Data gaps	8
	Exposure-response relationship	9
	Co-linearity	10
3	Previous research	12
	Mortality	12
	Hospitalisation	13
	Sydney	13
	Melbourne	15
	Brisbane	16
	Perth	17
	Multiple cities	17
4	Proposed methodology	19
	Effect estimate selection	20
	Data modifications to account for the unknown exposure-response relationship	21
	Notes on the hospital data analysis	22
5	Results	24
6	Conclusions and future challenges	26

Appendix A: Key pollutants	28
Particulate matter (PM)	28
Main sources	28
Main measuring techniques	29
Recent trends	29
Ozone (O ₃)	29
Main sources	29
Main measuring technique	30
Recent trends	30
Nitrogen dioxide (NO ₂)	30
Main sources	30
Main measuring technique	30
Recent trends	30
Carbon monoxide (CO)	31
Main sources	31
Main measuring technique	31
Recent trends	31
Sulfur dioxide (SO ₂)	31
Main sources	31
Main measuring technique	31
Recent trends	32
References	33

Acknowledgments

Authors of this report included Adrian Webster, Kasha Krysiak and Perri Timmins of the Australian Institute of Health and Welfare (AIHW). Valuable contributions were also made by Bruce Bacon (independent statistical expert), Vanessa Prescott (AIHW), Kuldeep Bhatia (AIHW), Sean Walsh (Environment Protection Authority, Victoria) and Guy Marks and Leanne Poulos (Australian Centre for Asthma Monitoring).

Funding and review was provided by the Australian Government Department of Health and Ageing.

Summary

Air pollution can trigger asthma and, in severe cases, has been associated with hospitalisation and death. How much of a problem is air pollution though? What proportion of asthma exacerbations are due to air pollution each year? How much does this vary with extreme events such as bushfires and dust storms? These are important questions that have so far proved difficult to answer.

Many improvements have been made to the quality and quantity of air pollution data over recent decades. So far, however, analysis of the contribution of air pollution to health conditions like asthma in Australia has been restricted to individual studies and investigations.

This paper discusses the challenges associated with this type of monitoring, outlines the work that has so far been done in Australia in this area and presents a method for estimating the contribution of air pollution to asthma hospitalisations. The method has been applied to a particular case study—Melbourne in 2006—to test its potential usefulness.

The adjusted results of the case study suggest that:

- about 3% of all asthma hospitalisations in Melbourne in 2006 were related to exposure to nitrogen dioxide
- about 4% of asthma hospitalisations of 0–14 year olds were related to particulates in the air.

The broader conclusion drawn is that the method used could provide a guide to monitoring the impact of air pollution on asthma exacerbations over time. The method requires refinement, however, and its application would be dependent on filling key gaps in the available research. If this method (or any similar method) is to play a role in the systematic monitoring of the health impacts of air pollution nationally, a targeted research programme is required that would provide:

- a national set of relative risk estimates that quantify the risks associated with exposure to key pollutants
- a more detailed understanding of the exposure-response relationship for each pollutant.