

Indicator 3a: Interval cancer rate

Table 19: Numbers and age-specific rates of interval cancers in women screened during 1996, 1997 and 1998, first screening round, 0-12 months, states and territories

Age group	Number / Rate	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40-49	Number	72	19	26	14	7	2	3	n.a.	n.a.
	Rate	8.4	5.4	5.2	6.1	5.3	3.4	8.4	n.a.	n.a.
50-59	Number	71	49	25	15	22	1	0	n.a.	n.a.
	Rate	7.7	7.2	5.0	7.0	12.3	2.0	0.0	n.a.	n.a.
60-69	Number	37	21	11	5	2	1	2	n.a.	n.a.
	Rate	6.8	7.1	3.9	4.9	2.8	12.7	15.8	n.a.	n.a.
70+	Number	20	11	4	1	0	1	0	n.a.	n.a.
	Rate	7.6	7.6	3.5	2.8	0.0	10.8	0.0	n.a.	n.a.
Ages 40+	Number	200	100	66	35	31	5	5	n.a.	n.a.
	Crude rate	7.7	6.8	4.7	6.0	7.5	3.5	5.5	n.a.	n.a.
	ASR (A)	7.6	6.9	4.5	5.7	6.7	3.9	6.1	n.a.	n.a.
	95% CI	6.4-8.7	5.5-8.1	3.3-5.7	3.8-7.9	4.3-9.5	0.7-8.3	0.6-13.6	n.a.	n.a.
Ages 50-69	Number	108	70	36	20	24	2	2	n.a.	n.a.
	Crude rate	7.4	7.2	4.6	6.3	9.6	2.6	4.1	n.a.	n.a.
	ASR (A)	7.3	7.2	4.5	6.1	8.4	2.8	6.6	n.a.	n.a.
	95% CI	5.9-8.8	5.6-8.9	3.1-6.1	3.5-9.0	4.8-12.1	0.0-7.4	0.0-16.5	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of interval cancers detected per 10,000 women-years and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. Northern Territory data were unavailable at the time of publication.
3. NSW rates include women with a personal history of breast cancer in the denominator.
4. The data include both symptomatic and asymptomatic women.

Source: AIHW analysis of BreastScreen Australia data.

Table 20: Numbers and age-specific rates of interval cancers in women screened during 1996, 1997 and 1998, first screening round, 13–24 months, states and territories

Age group	Number / Rate	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Number	n.a.	44	38	23	19	5	2	n.a.	n.a.
	Rate	n.a.	12.5	8.0	10.2	15.3	8.8	5.9	n.a.	n.a.
50–59	Number	n.a.	93	65	24	19	7	5	n.a.	n.a.
	Rate	n.a.	13.7	13.4	11.4	10.6	14.1	14.2	n.a.	n.a.
60–69	Number	n.a.	30	37	12	5	3	2	n.a.	n.a.
	Rate	n.a.	10.2	13.4	12.0	7.1	12.7	16.4	n.a.	n.a.
70+	Number	n.a.	17	14	4	1	0	0	n.a.	n.a.
	Rate	n.a.	11.8	12.4	11.6	3.1	0.0	0.0	n.a.	n.a.
Ages 40+	Number	n.a.	184	154	63	44	15	9	n.a.	n.a.
	Crude rate	n.a.	12.5	11.4	11.0	10.8	10.7	10.3	n.a.	n.a.
	ASR (A)	n.a.	12.2	12.2	11.3	9.7	10.9	11.3	n.a.	n.a.
	95% CI	n.a.	10.5–14.0	10.1–14.1	8.2–14.4	6.5–12.8	5.4–17.4	4.5–20.5	n.a.	n.a.
Ages 50–69	Number	n.a.	123	102	36	24	10	7	n.a.	n.a.
	Crude rate	n.a.	12.6	13.4	11.6	9.6	13.7	14.7	n.a.	n.a.
	ASR (A)	n.a.	12.2	13.4	11.6	9.2	13.5	15.1	n.a.	n.a.
	95% CI	n.a.	10.1–14.3	10.8–15.9	7.5–15.5	5.4–12.9	5.3–22.4	5.0–28.5	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of interval cancers detected per 10,000 women-years and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. New South Wales and Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Table 21: Numbers and age-specific rates of interval cancers in women screened during 1996, 1997 and 1998, first screening round, 0–24 months, states and territories

Age group	Number / Rate	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Number	n.a.	63	64	37	26	7	5	n.a.	n.a.
	Rate	n.a.	9.0	6.6	8.1	10.1	6.1	7.2	n.a.	n.a.
50–59	Number	n.a.	142	90	39	41	8	5	n.a.	n.a.
	Rate	n.a.	10.5	9.1	9.2	11.4	8.0	7.0	n.a.	n.a.
60–69	Number	n.a.	51	48	17	7	4	4	n.a.	n.a.
	Rate	n.a.	8.6	8.6	8.4	5.0	8.3	16.1	n.a.	n.a.
70+	Number	n.a.	28	18	5	1	1	0	n.a.	n.a.
	Rate	n.a.	9.7	7.9	7.2	1.5	5.0	0.0	n.a.	n.a.
Ages 40+	Number	n.a.	284	220	98	75	20	14	n.a.	n.a.
	Crude rate	n.a.	9.7	8.0	8.5	9.1	7.0	7.9	n.a.	n.a.
	ASR (A)	n.a.	9.5	8.3	8.5	8.1	7.3	8.7	n.a.	n.a.
	95% CI	n.a.	8.4–10.6	7.1–9.5	6.7–10.4	6.3–10.2	4.2–11.0	4.0–13.9	n.a.	n.a.
	Ages 50–69	Number	n.a.	193	138	56	48	12	9	n.a.
	Crude rate	n.a.	9.9	8.9	8.9	9.6	8.1	9.3	n.a.	n.a.
	ASR (A)	n.a.	9.7	8.9	8.9	8.8	8.1	10.8	n.a.	n.a.
	95% CI	n.a.	8.3–11.1	7.5–10.4	6.6–11.3	6.3–11.5	3.8–13.0	4.1–18.3	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of interval cancers detected per 10,000 women-years and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. New South Wales and Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Table 22: Numbers and age-specific rates of interval cancers in women screened during 1996, 1997 and 1998, subsequent screening rounds, 0–12 months, states and territories

Age group	Number / Rate	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Number	70	15	21	15	18	3	5	n.a.	n.a.
	Rate	8.6	9.6	4.3	9.3	12.6	3.3	9.1	n.a.	n.a.
50–59	Number	158	102	55	45	39	15	8	n.a.	n.a.
	Rate	9.0	7.5	7.3	8.5	7.5	9.0	7.8	n.a.	n.a.
60–69	Number	113	80	38	33	37	8	9	n.a.	n.a.
	Rate	7.5	6.6	6.7	8.0	8.0	6.1	14.8	n.a.	n.a.
70+	Number	48	22	18	2	2	2	1	n.a.	n.a.
	Rate	6.0	4.5	5.7	3.5	2.7	10.9	9.9	n.a.	n.a.
Ages 40+	Number	389	219	132	95	96	28	23	n.a.	n.a.
	Crude rate	8.0	6.8	6.2	8.2	8.0	6.9	10.0	n.a.	n.a.
	ASR (A)	8.1	7.3	6.3	7.9	8.1	7.2	10.3	n.a.	n.a.
	95% CI	7.3–9.0	6.1–8.5	5.2–7.4	6.2–9.6	6.4–9.8	4.3–10.4	6.2–14.8	n.a.	n.a.
	Ages 50–69	Number	271	182	93	78	76	23	17	n.a.
	Crude rate	8.3	7.1	7.0	8.3	7.7	7.7	10.4	n.a.	n.a.
	ASR (A)	8.4	7.1	7.0	8.3	7.7	7.8	10.7	n.a.	n.a.
	95% CI	7.4–9.3	6.1–8.3	5.6–8.5	6.5–10.1	6.0–9.4	4.5–11.1	5.8–16.6	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of interval cancers detected per 10,000 women-years and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. Northern Territory data were unavailable at the time of publication.
3. NSW rates include women with a personal history of breast cancer in the denominator.

Source: AIHW analysis of BreastScreen Australia data.

Table 23: Numbers and age-specific rates of interval cancers in women screened during 1996, 1997 and 1998, subsequent screening rounds, 13–24 months, states and territories

Age group	Number / Rate	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Number	n.a.	16	46	12	18	8	4	n.a.	n.a.
	Rate	n.a.	10.3	10.0	9.2	14.3	9.2	8.0	n.a.	n.a.
50–59	Number	n.a.	191	104	47	58	17	17	n.a.	n.a.
	Rate	n.a.	14.1	14.4	10.3	11.1	10.7	17.9	n.a.	n.a.
60–69	Number	n.a.	159	87	29	58	12	6	n.a.	n.a.
	Rate	n.a.	13.1	15.7	8.3	12.5	9.6	10.6	n.a.	n.a.
70+	Number	n.a.	55	36	3	9	1	2	n.a.	n.a.
	Rate	n.a.	11.3	11.6	6.5	12.1	5.9	20.3	n.a.	n.a.
Ages 40+	Number	n.a.	421	273	91	143	38	29	n.a.	n.a.
	Crude rate	n.a.	13.1	13.3	9.3	12.1	9.8	13.8	n.a.	n.a.
	ASR (A)	n.a.	12.7	13.5	9.1	12.3	9.5	14.1	n.a.	n.a.
	95% CI	n.a.	11.2–14.3	11.9–15.2	7.1–11.1	10.0–14.6	6.5–12.8	8.9–20.1	n.a.	n.a.
Ages 50–69	Number	n.a.	350	191	76	116	29	23	n.a.	n.a.
	Crude rate	n.a.	13.6	15.0	9.5	11.8	10.2	15.2	n.a.	n.a.
	ASR (A)	n.a.	13.7	15.0	9.5	11.7	10.3	14.9	n.a.	n.a.
	95% CI	n.a.	12.2–15.1	12.9–17.0	7.2–11.6	9.5–13.9	6.6–14.3	8.9–20.9	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of interval cancers detected per 10,000 women-years and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. New South Wales and Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Table 24: Numbers and age-specific rates of interval cancers in women screened during 1996, 1997 and 1998, subsequent screening rounds, 0–24 months, states and territories

Age group	Number / Rate	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Number	n.a.	31	67	27	36	11	9	n.a.	n.a.
	Rate	n.a.	9.9	7.1	9.2	13.4	6.2	8.6	n.a.	n.a.
50–59	Number	n.a.	293	159	92	97	32	25	n.a.	n.a.
	Rate	n.a.	10.8	10.8	9.3	9.3	9.8	12.6	n.a.	n.a.
60–69	Number	n.a.	239	125	62	95	20	15	n.a.	n.a.
	Rate	n.a.	9.9	11.2	8.1	10.2	7.8	12.8	n.a.	n.a.
70+	Number	n.a.	77	54	5	11	3	3	n.a.	n.a.
	Rate	n.a.	7.9	8.6	4.8	7.4	8.5	15.0	n.a.	n.a.
Ages 40+	Number	n.a.	640	405	186	239	66	52	n.a.	n.a.
	Crude rate	n.a.	10.0	9.7	8.7	10.0	8.3	11.8	n.a.	n.a.
	ASR (A)	n.a.	10.0	9.8	8.4	10.2	8.3	12.1	n.a.	n.a.
	95% CI	n.a.	9.0–11.0	8.9–10.8	7.1–9.7	8.7–11.6	6.2–10.4	8.5–15.7	n.a.	n.a.
Ages 50–69	Number	n.a.	532	284	154	192	52	40	n.a.	n.a.
	Crude rate	n.a.	10.4	10.9	8.8	9.7	9.0	12.7	n.a.	n.a.
	ASR (A)	n.a.	10.4	10.9	8.8	9.7	9.0	12.7	n.a.	n.a.
	95% CI	n.a.	9.5–11.3	9.6–12.2	7.4–10.2	8.3–11.0	6.4–11.4	8.7–16.6	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of interval cancers detected per 10,000 women-years and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. New South Wales and Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Indicator 3b: Program sensitivity

Table 25: Program sensitivity rates for women screened during 1996, 1997 and 1998, first screening round, 0–12 months, states and territories

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
	(Per cent)								
40–49	71.0	81.4	83.2	77.8	82.9	83.3	75.0	n.a.	n.a.
50–59	85.3	87.7	89.4	81.5	80.4	96.7	100.0	n.a.	n.a.
60–69	90.7	92.1	94.5	93.9	97.1	93.8	83.3	n.a.	n.a.
70+	93.9	94.7	97.6	97.6	100.0	91.7	100.0	n.a.	n.a.
Ages 40+									
Crude rate	86.3	89.7	91.3	86.9	88.4	92.9	90.0	n.a.	n.a.
ASR (A)	84.9	88.5	90.6	86.2	88.0	92.5	90.2	n.a.	n.a.
95% CI	83.0–86.7	86.1–90.5	88.5–92.7	81.7–90.1	83.8–91.6	84.9–98.3	81.3–97.7	n.a.	n.a.
Ages 50–69									
Crude rate	87.7	89.5	91.7	87.7	86.7	95.7	93.9	n.a.	n.a.
ASR (A)	87.5	89.5	91.5	86.6	87.3	95.5	93.1	n.a.	n.a.
95% CI	85.4–89.8	87.1–91.9	89.0–94.0	81.1–92.1	82.4–92.0	88.3–100.0	82.7–100.0	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of screen-detected cancers as a percentage of all cancers (screen-detected and interval cancers) and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Table 26: Program sensitivity rates for women screened during 1996, 1997 and 1998, first screening round, 0–24 months, states and territories

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
(Per cent)									
40–49	n.a.	56.8	66.8	68.1	57.6	62.5	64.3	n.a.	n.a.
50–59	n.a.	71.1	70.1	73.3	72.0	78.4	80.8	n.a.	n.a.
60–69	n.a.	82.7	79.7	85.6	90.4	78.9	71.4	n.a.	n.a.
70+	n.a.	87.5	90.2	97.6	97.9	91.7	100.0	n.a.	n.a.
Ages 40+									
Crude rate	n.a.	75.5	75.9	79.2	77.7	77.4	76.3	n.a.	n.a.
ASR (A)	n.a.	73.4	74.6	78.7	77.4	76.9	77.2	n.a.	n.a.
95% CI	n.a.	70.6–76.0	71.7–77.6	73.6–83.3	72.5–81.5	67.9–84.9	66.3–87.0	n.a.	n.a.
Ages 50–69									
Crude rate	n.a.	75.5	74.3	79.4	78.8	78.6	77.5	n.a.	n.a.
ASR (A)	n.a.	75.9	74.1	78.4	79.7	78.6	76.9	n.a.	n.a.
95% CI	n.a.	72.9–78.9	70.6–77.8	71.8–85.0	74.1–85.1	66.7–88.3	62.8–89.6	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of screen-detected cancers as a percentage of all cancers (screen-detected and interval cancers) and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. New South Wales and Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Table 27: Program sensitivity rates for women screened during 1996, 1997 and 1998, subsequent screening rounds, 0–12 months, states and territories

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
(Per cent)									
40–49	63.9	70.0	77.9	65.1	63.3	84.2	66.7	n.a.	n.a.
50–59	78.0	81.0	79.2	77.8	82.4	72.7	80.5	n.a.	n.a.
60–69	84.6	87.0	85.2	84.9	85.0	66.7	69.0	n.a.	n.a.
70+	89.5	92.4	90.9	94.4	63.3	86.7	80.0	n.a.	n.a.
Ages 40+									
Crude rate	81.5	85.3	83.8	81.0	79.9	73.4	74.4	n.a.	n.a.
ASR (A)	78.4	81.8	82.1	79.2	76.8	75.2	74.4	n.a.	n.a.
95% CI	76.2–80.3	78.6–84.9	79.2–85.0	74.9–83.1	73.1–80.8	67.4–82.2	62.8–89.6	n.a.	n.a.
Ages 50–69									
Crude rate	81.4	84.2	82.2	81.5	83.8	69.7	75.7	n.a.	n.a.
ASR (A)	80.8	83.5	81.7	80.8	83.5	70.2	75.7	n.a.	n.a.
95% CI	78.7–83.0	81.2–85.7	78.3–85.0	76.7–84.7	79.8–86.8	60.8–79.2	64.9–83.3	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of screen-detected cancers as a percentage of all cancers (screen-detected and interval cancers) and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Table 28: Program sensitivity rates for women screened during 1996, 1997 and 1998, subsequent screening rounds, 0–24 months, states and territories

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
(Per cent)									
40–49	n.a.	53.0	52.5	65.1	46.3	59.3	52.6	n.a.	n.a.
50–59	n.a.	59.8	56.9	77.8	65.2	55.6	56.9	n.a.	n.a.
60–69	n.a.	69.1	63.7	84.9	68.9	69.7	57.1	n.a.	n.a.
70+	n.a.	77.7	76.8	94.4	84.1	81.3	57.1	n.a.	n.a.
Ages 40+									
Crude rate	n.a.	66.6	62.7	81.7	66.8	63.5	56.3	n.a.	n.a.
ASR (A)	n.a.	63.2	60.3	80.1	64.6	63.5	56.1	n.a.	n.a.
95% CI	n.a.	60.1–66.2	57.2–63.5	75.7–84.0	60.9–68.4	55.9–70.4	46.8–65.5	n.a.	n.a.
Ages 50–69									
Crude rate	n.a.	64.6	60.2	81.9	67.1	62.3	57.0	n.a.	n.a.
ASR (A)	n.a.	63.7	59.7	81.0	66.7	61.4	57.0	n.a.	n.a.
95% CI	n.a.	61.1–66.1	56.2–63.3	77.2–84.9	62.9–70.7	53.1–69.5	46.7–67.6	n.a.	n.a.

n.a. Not available.

Notes

1. Rates are the number of screen-detected cancers as a percentage of all cancers (screen-detected and interval cancers) and age-standardised to the population of women attending a BreastScreen Australia service in 1998.
2. New South Wales and Northern Territory data were unavailable at the time of publication.

Source: AIHW analysis of BreastScreen Australia data.

Indicator 4: Ductal carcinoma in situ

Table 29a: Numbers of women screened and cases of DCIS detected in these women, 1999, by age, states and territories

Age group Number		NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Screened	59,392	17,395	39,041	12,083	10,019	4,364	3,007	1,099	146,400
	Cases	31	14	20	12	11	2	4	0	94
50–59	Screened	97,064	78,163	55,598	26,016	28,578	8,040	5,637	2,001	301,097
	Cases	78	69	43	31	24	17	9	0	271
60–69	Screened	72,693	51,612	37,510	18,641	20,923	5,990	3,098	702	211,169
	Cases	75	53	52	18	21	6	2	0	227
70+	Screened	44,846	24,196	21,679	3,254	4,607	983	513	204	100,282
	Cases	47	21	24	5	10	2	2	0	111
Ages 40+	Screened	273,995	171,366	153,828	59,994	64,127	19,377	12,255	4,006	758,948
	Cases	231	157	139	66	66	27	17	0	703
Ages 50–69	Screened	169,757	129,775	93,108	44,657	49,501	14,030	8,735	2,703	512,266
	Cases	153	122	95	49	45	23	11	0	498

Source: AIHW analysis of BreastScreen Australia data.

Table 29b: Numbers of women screened and cases of DCIS detected in these women, 2000, by age, states and territories

Age group Number		NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
40–49	Screened	59,801	16,290	40,669	11,833	10,646	4,839	2,816	1,138	148,032
	Cases	44	25	19	10	6	3	4	1	112
50–59	Screened	98,467	78,018	61,673	29,495	29,517	9,105	5,541	2,127	313,943
	Cases	90	77	62	44	44	11	7	1	336
60–69	Screened	72,328	55,572	39,584	20,500	20,114	6,310	2,664	635	217,707
	Cases	63	60	44	33	25	8	3	1	237
70+	Screened	46,863	27,352	21,803	3,753	5,216	1,060	442	247	106,736
	Cases	55	39	29	7	13	1	1	0	145
Ages 40+	Screened	277,459	177,232	163,729	65,581	65,493	21,314	11,463	4,147	786,418
	Cases	252	201	154	94	88	23	15	3	830
Ages 50–69	Screened	170,795	133,590	101,257	49,995	49,631	15,415	8,205	2,762	531,650
	Cases	153	137	106	77	69	19	10	2	573

Source: AIHW analysis of BreastScreen Australia data.

Table 30a: Age-specific rates of DCIS detected in women screened, 1999, states and territories

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
(Per cent)									
40–49	5.2	8.0	5.1	9.9	11.0	4.6	13.3	0.0	6.4
50–59	8.0	8.8	7.7	11.9	8.4	21.1	16.0	0.0	9.0
60–69	10.3	10.3	13.9	9.7	10.0	10.0	6.5	0.0	10.7
70+	10.5	8.7	11.1	15.4	21.7	20.3	39.0	0.0	11.1
Ages 40+									
Crude rate	8.4	9.2	9.0	11.0	10.3	13.9	13.9	0.0	9.3
ASR (A)	8.4	9.0	9.3	11.3	11.0	14.5	15.6	..	9.2
95% CI	7.2–9.5	7.5–10.5	7.9–10.8	8.5–14.3	8.2–13.8	8.9–20.6	8.0–24.8	..	8.5–9.9
Ages 50–69									
Crude rate	9.0	9.4	10.2	11.0	9.1	16.4	12.6	0.0	9.7
ASR (A)	9.0	9.4	10.3	11.0	9.1	16.5	12.0	..	9.7
95% CI	7.5–10.4	7.6–11.1	8.2–12.4	8.1–13.9	6.5–11.7	10.0–23.1	5.5–19.9	..	8.8–10.6

.. Not applicable.

Note: Rates are the number of cases of DCIS per 10,000 women screened and age-standardised to the population of women attending a BreastScreen Australia service in 1998.

Source: AIHW analysis of BreastScreen Australia data.

Table 30b: Age-specific rates of DCIS detected in women screened, 2000, states and territories

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
(Per cent)									
40–49	7.4	15.3	4.7	8.5	5.6	6.2	14.2	8.8	7.6
50–59	9.1	9.9	10.1	14.9	14.9	12.1	12.6	4.7	10.7
60–69	8.7	10.8	11.1	16.1	12.4	12.7	11.3	15.7	10.9
70+	11.7	14.3	13.3	18.7	24.9	9.4	22.6	0.0	13.6
Ages 40+									
Crude rate	9.1	11.3	9.4	14.3	13.4	10.8	13.1	7.2	10.6
ASR (A)	8.9	11.8	9.6	14.4	13.5	10.7	13.8	8.0	10.5
95% CI	7.8–10.1	10.0–13.5	8.1–11.2	11.4–17.3	10.5–16.6	6.3–15.3	6.7–22.3	0.0–18.6	9.7–11.2
Ages 50–69									
Crude rate	9.0	10.3	10.5	15.4	13.9	12.3	12.2	7.2	10.8
ASR (A)	8.9	10.3	10.5	15.4	13.9	12.3	12.1	9.3	10.8
95% CI	7.4–10.4	8.5–12.0	8.4–12.5	12.0–18.8	10.7–17.1	6.5–18.2	5.3–19.9	0.0–26.2	9.9–11.7

Note: Rates are the number of cases of DCIS per 10,000 women screened and age-standardised to the population of women attending a BreastScreen Australia service in 1998.

Source: AIHW analysis of BreastScreen Australia data.