



**Australian Government**

**Australian Institute of  
Health and Welfare**

**AIHW**

# Hearing health outreach services for Aboriginal and Torres Strait Islander children in the Northern Territory

July 2012 to  
December 2019





# Hearing health outreach services for Aboriginal and Torres Strait Islander children in the Northern Territory

July 2012 to  
December 2019



The Australian Institute of Health and Welfare is a major national agency whose purpose is to create authoritative and accessible information and statistics that inform decisions and improve the health and welfare of all Australians.

© Australian Institute of Health and Welfare 2020



This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC-BY 3.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at [www.aihw.gov.au/copyright/](http://www.aihw.gov.au/copyright/). The full terms and conditions of this licence are available at [creativecommons.org/licenses/by/3.0/au/](http://creativecommons.org/licenses/by/3.0/au/).

ISBN 978-1-76054-709-7 (Online)

ISBN 978-1-76054-710-3 (Print)

### Suggested citation

Australian Institute of Health and Welfare 2020. Hearing health outreach services for Aboriginal and Torres Strait Islander children in the Northern Territory: July 2012 to December 2019. Cat no. IHW 228. Canberra: AIHW.

### Australian Institute of Health and Welfare

Board Chair  
Mrs Louise Markus


Chief Executive Officer  
Mr Barry Sandison

Any enquiries relating to copyright or comments on this publication should be directed to:  
Australian Institute of Health and Welfare  
GPO Box 570  
Canberra ACT 2601  
Tel: (02) 6244 1000  
Email: [info@aihw.gov.au](mailto:info@aihw.gov.au)

Published by the Australian Institute of Health and Welfare.

Please note that there is the potential for minor revisions of data in this report.  
Please check the online version at [www.aihw.gov.au](http://www.aihw.gov.au) for any amendments.





# Contents

Summary.....	v
<b>1 Introduction.....</b>	<b>1</b>
Why is ear and hearing health important? .....	2
What is ear and hearing health? .....	2
Impact of poor ear and hearing health .....	2
Ear and hearing health in the Northern Territory.....	2
Australian Government-funded Hearing Health Program in the Northern Territory.....	3
How do children and young people move through the NT hearing health system? .....	4
About this report.....	5
About the data in this report .....	5
<b>2 Service delivery.....</b>	<b>6</b>
Health education, promotion and prevention.....	7
Outreach audiology .....	8
ENT teleotology services .....	10
Clinical Nurse Specialist services.....	12
<b>3 Ear conditions and hearing health status .....</b>	<b>14</b>
Ear conditions.....	15
Age and sex differences .....	16
Changes among children and young people who received 2 or more services between 2012 and 2019.....	18
Ear conditions among CNS service recipients .....	19
Hearing status .....	20
Hearing loss .....	21
Variation by age and sex .....	21
Hearing loss among children and young people with ear conditions.....	22
Trends.....	22
Hearing impairment.....	23
Variation by age and sex .....	23
Trends.....	24
Changes over time in hearing loss and impairment.....	25



<b>4 Demand for ear and hearing health services and other follow-up services</b> .....	<b>27</b>
Follow-up services required after audiologist visits .....	28
Follow-up services required after ENT teleotology services .....	28
Waiting list for audiology services and ENT teleotology services .....	29
<b>5 Regional analysis</b> .....	<b>30</b>
Hearing Loss .....	31
Hearing Impairment .....	33
Ear Conditions .....	35
<b>6 Progress against benchmarks</b> .....	<b>37</b>
Service delivery .....	38
Health outcomes—hearing impairment .....	39
Health outcomes—middle ear conditions .....	39
<b>Appendix A: About the Hearing Health Program data collections</b> .....	<b>40</b>
Data collection, management and reporting .....	40
<b>Appendix B: Data quality statement</b> .....	<b>41</b>
Outreach audiology data collection summary .....	41
ENT teleotology data collection summary .....	41
CNS data collection summary .....	42
Child Health Check Initiative (CHCI) data collections summary .....	42
<b>Acknowledgments</b> .....	<b>43</b>
<b>Abbreviations</b> .....	<b>44</b>
<b>Symbols</b> .....	<b>44</b>
<b>Glossary</b> .....	<b>45</b>
<b>References</b> .....	<b>47</b>
<b>List of figures</b> .....	<b>48</b>
<b>List of tables</b> .....	<b>49</b>
<b>List of boxes</b> .....	<b>49</b>
<b>Related publications</b> .....	<b>50</b>



## Summary

Ear and hearing health is vital for overall health and quality of life. Ear disease and associated hearing loss can have long-lasting impacts on education, wellbeing and employment.

Aboriginal and Torres Strait Islander children are more likely than non-Indigenous children to experience ear and hearing problems. A number of factors contribute to the poorer ear and hearing health of Indigenous children, including lack of access to health services, household overcrowding and second-hand smoke exposure. Since 2018, the Australian Government has funded the Northern Territory Government to deliver hearing health outreach services available to more than 31,500 Indigenous children and young people aged under 21 in the Northern Territory.

This report presents new data for the Hearing Health outreach services for 2019. It focuses on services provided in 2019 and produces time trends between July 2012 and December 2019. Throughout this report, links to the supplementary tables relate to the figures presented. Supplementary tables are available at the AIHW website.

### How many children received services?

- In 2019, 137 training sessions for hearing health education, promotion and prevention activities were provided to health-care staff.
- A total of 1,921 children and young people received at least 1 audiology, Clinical Nurse Specialist (CNS) or ear, nose and throat (ENT) teleotology service in 2019. From July 2012 to December 2019, a total of 29,930 services were provided to more than 8,000 children and young people, an average of 3 services per child and young person.
- In 2019, 2,156 outreach audiology services were provided to 1,896 children and young people. From July 2012 to December 2019, a total of 16,524 services were provided to a total of 7,899 children and young people, an average of 2 services per unique service recipient.
- 770 ENT teleotology services were provided to 709 children and young people. From July 2012 to December 2019, a total of 6,841 services were provided to a total of 3,609 children and young people, an average of 2 services per unique service recipient.
- CNSs conducted 1,119 visits to 1,008 children. From July 2012 to December 2019, a total of 6,515 services were provided to a total of 4,521 children, an average of about 1 visits per unique service recipient.

**Figure 1: Number of children and young people who received audiology, CNS or ENT teletology service in 2019**



**1,896**

children and young people received **audiology** services



**1,008**

children and young people received **CNS** services



**709**

children and young people received **ENT teletology** services

Sources: tables S2.1, S2.3 and S2.5.

## Hearing health is improving among children and young people who have received hearing health outreach services between 2012 and 2019

The percentage of children and young people with at least 1 ear disease decreased by 4 percentage points, from 66% (483 among 727 children and young people) to 62% (1,188 among 1,921 children and young people).

The percentage of children and young people with hearing loss decreased by 13 percentage points, from 55% (388 of 791 children and young people) to 42% (793 of 1,893 children and young people).

**Figure 2: Ear and hearing health improvements between 2012 and 2019:**

**66%**



**62%**

Children and young people with at least 1 **ear disease**



**55%**



**42%**

Children and young people with **hearing loss**

Sources: tables S3.1 and S3.6.

Among 3,480 children who received at least 2 services between 2012 and 2019 and who had hearing loss at their first service, more than 1,550 (61%) had improved—a change from bilateral to unilateral hearing loss or from unilateral hearing loss to no hearing loss.

Among 1,647 children who had hearing impairment at their first service, 1,165 (71%) had improved—moving to a lower degree of hearing impairment or to no hearing impairment.

## Many children and young people are still waiting for hearing health services

At December 2019, there were 3,459 children and young people on the audiology waiting list. Among them, 1,726 were new referrals and 1,733 were recalled for further follow-up after receiving an initial audiology service. There were 2,458 children and young people on the waiting list for ENT teleotology services. Among them, 1,025 were new referrals and 1,433 were recalled for further follow-up after receiving initial ENT teleotology services.

### Is the Hearing Health Program meeting its benchmarks?

The annual benchmark for hearing health service delivery is measured by the number of services per year. From 2012 to 2019, service delivery targets for audiology services, CNS services and health promotion services were either met or exceeded.

Health outcomes in this program are measured every 3 years (July 2012–June 2015, July 2015–December 2018 and January 2019–June 2021). All benchmarks in the first 2 periods were met. As 2019 is the first year of the new period, the information in this report is based on the data from 1 year only and does not indicate whether the benchmarks have been met or not (Table 1).


The progress against benchmarks in 2019 is presented in Table 1.

**Table 1: Progress against benchmarks, 2019**

2019–2021 Target	Outcome in 2019
At least 1,700 audiology services per year	2,156 audiology services provided
At least 700 children receiving CNS services per year	1,008 children received CNS services
At least 60 hearing health promotion or training services/activities per year	137 preventative hearing health promotion or training services and activities provided in 2019
Less than 7.5% of children tested with moderate/severe conductive hearing impairment during 2019–2021	4% of children tested with moderate/severe conductive hearing impairment, January–Dec 2019
Less than 7% of children (0–5) with CSOM during 2019–2021	8% of children (0–5) with CSOM, January–Dec 2019
Less than 7% of children (6–15) with CSOM during 2019–2021	7% of children (6–15) with CSOM, January–Dec 2019
Less than 5% of children (0–5) with dry perforation 2019–2021	7% of children (0–5) with dry perforation, January–Dec 2019
Less than 9% of children (6–15) with dry perforation 2019–2021	16% of children (6–15) with dry perforation, January–Dec 2019

CSOM = Chronic Suppurative Otitis Media





# 1

## Introduction



## Why is ear and hearing health important?

Hearing loss is more prevalent among Aboriginal and Torres Strait Islander people than among non-Indigenous Australians, and continues to be an important health and social issue. Compared with non-Indigenous children, Indigenous children are reported to have 2.9 times the rate of ear and hearing problems (AIHW 2017a).

### What is ear and hearing health?

'Ear and hearing health' can refer to a variety of ear-related conditions: hearing impairment, hearing loss, and the relationship between these health problems. 'Hearing loss' involves loss of hearing in 1 or both ears, and 'hearing impairment' describes the degree of impairment associated with hearing loss in the 'better hearing ear'.

'Middle ear disease' includes conditions such as otitis media and its various forms. Otitis media refers to all forms of inflammation of the middle ear, and can commonly be caused by infection or Eustachian tube dysfunction (ETD). The Eustachian tube functions to equalise the pressure in the middle ear to atmospheric pressure and impairment of this function is known as ETD. Eustachian tube function can be impaired for a number of reasons, most commonly occurring in childhood when the Eustachian tube is developing. Enlarged adenoids (glands in the roof of the mouth) are often present.

### Impact of poor ear and hearing health

Among Indigenous children, otitis media is a large contributor to hearing loss, and often manifests itself at earlier ages, with greater severity, greater persistence and more frequently than in non-Indigenous children (Jervis-Bardy et al. 2014).

Hearing loss can also have severe negative impacts on language development, cognitive development and socialisation, particularly in infants and young children. Hearing loss in early childhood can lead to social, learning, linguistic and behavioural problems in school. Experiencing these difficulties can translate into a lifetime of disadvantage, affecting areas such as wellbeing, social success, income, and employment (WHO 1996).

## Ear and hearing health in the Northern Territory

Middle ear disease is a common health problem and is a cause for concern among Indigenous children, particularly those who live in remote communities (ABS 2016). This may be associated with several factors, including:

- the nature of otitis media, which makes it a complicated disease to manage
- the acceptance in many communities of otitis media as normal because it is common
- living conditions in some parts of the Northern Territory—for example, poorly maintained housing, household overcrowding and related hygiene issues
- second-hand smoke exposure



- the inability to find children and their families in communities during health outreach visits because of the high mobility of Indigenous families
- the geographical location and vast spread of Indigenous communities, which makes access to services difficult
- the difficulty in recruiting and retaining a specialist workforce
- the difficulty of the health sector to improve systemic problems with culturally appropriate health care and unconscious bias.

The last 2 factors are also associated with living in remote areas. Living in remote areas may also affect access to general and ear health services: there are fewer medical practitioners per capita, and less access to general practitioners (AIHW 2014a, 2014b). Decreased access to these services can result in delays in diagnosis, treatment and management of middle ear disease among Indigenous children, prolonging periods of hearing loss and impairment.

## Australian Government-funded Hearing Health Program in the Northern Territory

Hearing health services were expanded in the Northern Territory in response to the Child Health Check Initiative (CHCI) in July 2007, which was introduced under the Northern Territory Emergency Response (NTER). The original Child Health Check data for the 9,373 Aboriginal and Torres Strait Islander children who received services in the NTER Prescribed Areas showed that between July 2007 and June 2009, 30% had ear disease. Through the introduction of the CHCI, the children who were found to have ear diseases were able to obtain audiology, and ear, nose and throat (ENT) specialist services (AIHW & DoHA 2009).

The program continued under the Closing the Gap initiative (CtG) in the Northern Territory National Partnership Agreement from mid-2009 to mid-2012. More information can be found in the 2012 AIHW publication *Northern Territory Emergency Response Child Health Check Initiative—follow-up services for oral and ear health: final report 2007–2012* (AIHW 2012).

Between July 2012 and June 2015, the ear and hearing health services were replaced and expanded by the National Partnership Agreement on Stronger Futures in the Northern Territory (SFNT). Since July 2015, these services have been continued through the new National Partnership on Northern Territory Remote Aboriginal Investment (NTRAI).

The Australian Government also funds the Northern Territory Government to deliver services through the Healthy Ears—Better Hearing, Better Listening Program. The Northern Territory Government uses this funding to support ENT teleotology services that are part of the hearing health outreach services. The teleotology services are for children and young people aged under 21, especially in remote areas where there is high demand and lack of local services. The hearing health outreach services mentioned throughout this report refers to all of the Australian Government-funded programs that stemmed from the CHCI in the Northern Territory since July 2007.


There are 4 services in the Hearing Health Program (detailed information on these services can be found in Chapter 2):

1. health education, promotion and prevention
2. outreach audiology
3. ENT teleotology
4. Clinical Nurse Specialists (CNSs)—Case Management and Health Promotion.

## How do children and young people move through the Northern Territory hearing health system?

The hearing health outreach services are available to more than 31,510 Indigenous children and young people aged under 21 in the Northern Territory. As illustrated in Figure 1.1 below, children and young people generally enter the Northern Territory public hearing health system through the primary health-care sector. From this starting point, referrals can be made to audiologists, ENT specialists or CNS services. CNS services can also provide referrals to audiologists, or to other organisations. After children have been seen by audiologists or ENT specialists, they can be sent for follow-up in primary care, referred to other community-based support organisations, or referred to visiting rehabilitation support services through Hearing Australia or the Department of Education and Training’s Hearing Support Services.

**Figure 1.1: Clinical path of children and young people moving through the Northern Territory public hearing health system**





## About this report


This report presents information from July 2012 to December 2019 on hearing health outreach services provided to the Aboriginal and Torres Strait Islander children and young people aged under 21 in the Northern Territory. The Australian Government funds these services via various programs, and the Northern Territory Government delivers them. These programs aim to provide outreach services for the early detection, treatment and management of ear diseases and hearing health problems among Indigenous children and young people.

This report is an update of *Hearing health outreach services for Aboriginal and Torres Strait Islander children in the Northern Territory July 2012 to December 2018*, which is part of the Australian Institute of Health and Welfare's (AIHW) publication series reporting on the hearing programs in the Northern Territory. Throughout this report, links to the supplementary tables relate to the figures presented. Supplementary tables are available at: <https://www.aihw.gov.au/reports/adoption/hearing-services-for-aboriginal-and-tsi-children/data>.

## About the data in this report

The data used in this report are collected from the hearing health outreach services funded by the Department of Health. The data refer to more than 8,000 children and young people aged under 21 who received services between July 2012 and December 2019. This accounts for approximately 26% of the Northern Territory Indigenous population in this age group.

However, children and young people who received these services are not a random sample of the population. Audiology services were provided to children and young people in remote communities during visits from audiologists and specialists. Since January 2013, children and young people have been prioritised according to their need for services, which means that those with worse ear and hearing health are more likely to be captured in the data collection.



# 2

## Service delivery


## Health education, promotion and prevention

A variety of hearing health education, promotion and prevention activities are delivered through the Northern Territory public hearing health system. Priority areas of promotion and prevention centre on enhancing hearing health literacy through education and community participation. Health education, promotion and prevention initiatives include:

- training for Aboriginal community hearing workers—including training to prevent the first infection
- ear health promotion material such as handouts, posters and audiovisual messages
- a health promotion hip-hop music video addressing ear health
- development of 2 health promotion videos with key ear and hearing health messages and medication compliance.

In 2019, 137 training sessions for hearing health education, promotion and prevention activities were provided to health-care staff. The target audiences for these activities are illustrated below.

**Figure 2.1: Ear health education, promotion and prevention training sessions delivered in 2019:**






## Outreach audiology

Audiology services include assessing middle ear function, diagnosing hearing loss and middle ear disease, and recommending clinical care or rehabilitation (such as communication strategies, classroom amplification, hearing aids, speech therapy and educational support). These services are delivered by audiology outreach teams, which consist of an audiologist and at least 1 other member of staff such as a registered nurse, nurse audiometrist, or Aboriginal health practitioner.

Parents or guardians of service recipients must consent to sharing information with the AIHW. The demographic information in this report, apart from the number of services and service recipients, represents only children who have provided consent to share their information. When a child’s parent or guardian does not provide consent to share information, only a limited amount of aggregate information is provided to AIHW. See Appendix B for more information.

- In 2019, 2,156 audiology services were provided to 1,896 service recipients (Figure 2.2).
- From July 2012 to December 2019, a total of 16,524 services were provided to a total of 7,899 unique service recipients.
- The numbers of audiology services and service recipients increased slightly between 2018 and 2019.

**Figure 2.2: Number of outreach audiology services and service recipients, July 2012–December 2019:**



*Note:* The total number of service recipients in July 2012 to 2019 combined does not sum to the years because some children and young people received services in multiple calendar years; these services recipients were counted only once in the total.

*Source:* Table S2.1.


The drop in numbers between 2015 and 2019 was caused mainly by:

- difficulty locating children in the community due to transient families
- lack of community support to locate and transfer clients to the health centre.


ENT teleotology and CNS services experienced similar trends, with CNS services affected also by high staff turnover.

Overall, rates of non-consent to share information for audiology services are low, and in 2019 there were no cases of non-consent for audiology service recipients.

In 2019, more boys than girls received audiology services (Figure 2.3):



The highest proportion (34%) of service recipients in 2019 were aged 6–10 (Figure 2.4).



More information on outreach audiology services, such as services recommended by audiologists, can be found in the supplementary tables online.


## ENT teleotology services

The ENT teleotology service model was developed to meet the demand for ENT services in remote Northern Territory communities. An outreach visit from an audiologist and ENT nurse provides both hearing and full clinical assessments. Video-otoscopy is performed and video sequences are stored and forwarded to the ENT team at the Royal Darwin Hospital and Alice Springs Hospital. Using the video sequences and clinical data, the ENT specialist provides advice, diagnosis, assessment and treatment recommendations (for example, medications, care coordination, surgery, hearing aids).

The availability of teleotology services in remote communities ensures that services are culturally safe and cost-effective, and eliminates time that families would otherwise be away from the community for specialist appointments. These services also increase access to ENT services for children and reduce the burden on tertiary centres where demand for ENT services is high.

- In 2019, 770 ENT teleotology services were provided to 709 service recipients (Figure 2.5).
- From July 2012 to December 2019, a total of 6,841 services were provided to a total of 3,609 unique service recipients.
- Consent rates to share the information have been very high, and between 2016 and 2019 there were no instances of non-consent.

**Figure 2.5: Number of ENT teleotology services and service recipients, July 2012–December 2019**




*Note:* The total number of service recipients in July 2012 to 2019 combined does not sum to the years because some children and young people received services in multiple calendar years; these services recipients were counted only once in the total.

*Source:* Table S2.3.




The proportion of service recipients was slightly higher for boys than for girls in 2019 (Figure 2.6).

**Figure 2.6: ENT Teleotology service recipients in 2019, by sex**



The highest proportion (38%) of service recipients in 2019 were aged 6–10 (Figure 2.7).

**Figure 2.7: ENT Teleotology service recipients in 2019, by age group**



Source: Table S2.4.

More information on ENT services, such as services recommended by ENT specialists, can be found in the supplementary tables online.


## Clinical Nurse Specialist services

CNS services were developed in response to the challenges encountered in preventing ear disease and implementing clinical care for otitis media in the Northern Territory. The CNS is responsible for early identification and case management of high priority 0–5 year old children. The CNS oversees and coordinates the treatment, management and coordination of children with a prioritised need for care by acting as a central point of contact between primary health-care services and specialist resources.

In 2019, 1,119 CNS services were provided to 1,008 service recipients (Figure 2.8).

- From July 2012 to December 2019, a total of 6,515 services have been provided to a total of 4,521 unique service recipients.
- From 2012 to 2016, there was a substantial increase in the number of service recipients and services provided, followed by a decrease in 2017 and 2018, then increasing again in 2019.

**Figure 2.8: Number of CNS services and service recipients, July 2012–December 2019**




*Note:* The total number of service recipients in July 2012 to 2019 combined does not sum to the years because some children and young people received services in multiple calendar years; these services recipients were counted only once in the total.

*Source:* Table S2.5.

Caution should be taken when interpreting CNS data in 2013 and 2014 because non-consent rates for sharing information were high. However, non-consent rates have been decreasing over time. In 2015, the hearing health team in the Northern Territory Department of Health implemented a training program to improve consent rates, and this could explain the decrease in non-consent rates in recent years.


More boys than girls received CNS services in 2019 (Figure 2.9).

**Figure 2.9: Clinical Nurse Specialist service recipients in 2019, by sex**



The CNS services are available to Aboriginal and Torres Strait Islander children who have been assigned a hearing health priority (Figure 2.10)—priority is closely linked to age (Box 2.1).

**Figure 2.10: Hearing health priority (closely linked to age) among CNS service recipients, 2019**



Source: Table S2.6.

Older children represented the majority of service recipients in 2019: about 4 in 5 children were aged 3–10, and younger age groups represented a smaller proportion of service recipients.

More information on CNS services can be found in the supplementary tables (S2.5 and S2.6).


**Box 2.1: Hearing health priority (HP) categories**

**HP1:** Infants <12 months with recurrent acute otitis media (AOM) or chronic suppurative otitis media (CSOM). Infants who have failed newborn hearing screening.

**HP2:** Children aged 1–2 with perforation of the eardrum, recurrent AOM or persistent bilateral otitis media with effusion (OME).

**HP3:** Children aged 3–5 with perforation of the eardrum, recurrent AOM, persistent bilateral OME or moderate to profound hearing impairment.

**HP4:** Children aged 6–10 with moderate, severe or profound hearing impairment.



# 3

## **Ear conditions and hearing health status**

## Ear conditions

Two main types of ear conditions are captured in the Hearing Health Program: otitis media, and Eustachian tube dysfunction (ETD) (Box 3.1).

### Box 3.1: Types of ear conditions

**Otitis media:** all forms of inflammation and infection of the middle ear. Active inflammation or infection is nearly always associated with a middle ear effusion (fluid in the middle ear space). Types of otitis media include:


- **acute otitis media (AOM)**—the presence of fluid behind the eardrum plus at least 1 of the following: bulging eardrum, red eardrum, recent discharge of pus, fever, ear pain or irritability for less than 6 weeks
- **chronic suppurative otitis media (CSOM) with discharge**—a persistent suppurative discharge from the middle ear through a tympanic membrane perforation, for more than 6 weeks
- **chronic suppurative otitis media (CSOM) without discharge**—the presence of a perforation (hole) in the eardrum without evidence of discharge or fluid behind the eardrum (also known as inactive CSOM or dry perforation)
- **otitis media with effusion (OME)**—the presence of an intact eardrum and middle ear fluid without symptoms or signs of acute infection. OME may be episodic or persistent.

**Eustachian tube dysfunction (ETD):** negative middle ear pressure associated with compromised equalisation, impeding middle ear function and sometimes causing middle ear fluid accumulation.

In 2019, 1,921 children and young people received at least 1 audiology, CNS or ENT teleotology service. At their latest service, 1,188 children and young people were diagnosed with at least 1 type of ear condition. The proportion of children who were diagnosed with an ear condition has declined from 66% in 2012 to 62% in 2019, although the proportions of specific ear conditions have had small fluctuations over the years.

In 2019, the most common type of ear condition, among the 1,188 children diagnosed with an ear condition, was OME (25%, or 474 cases), followed by ETD (17%, or 321 cases); CSOM without discharge (15%, or 279 cases); and CSOM with discharge (9.3%, or 179 cases) (Figure 3.1).

**Figure 3.1: Types of ear conditions among children and young people who received an audiology, CNS or ENT teleotology service in 2019**



(a) 'Other' includes grommets, myringitis, otitis external, tympanosclerosis, and other ear conditions.

Source: Table S3.1.



**3 in 5 children  
(62%) were  
found to have  
≥1 ear condition**

### Age and sex differences


The proportions of ear conditions varied among age groups in 2019.

Younger children who received a service had the highest proportion of ear conditions. In 2019, 73% of children aged 0–2 and 56% of children aged 16–20 had at least 1 type of ear condition. This pattern reflects the natural profile of ear disease, where children typically grow out of ear conditions (AIHW 2014c).



The most common types of ear conditions differed among age groups in 2019, as shown in Figure 3.2. Similar patterns among age groups were seen in previous years.

**Figure 3.2: Most common ear conditions in children and young people who received at least 1 audiology outreach, CNS or ENT teleotology service in 2019, by age group**



Source: Table S3.2.

Apart from the trends shown above, the following patterns were also observed:

- The proportion of OME, ETD and AOM generally decreased with age.
- The proportion of CSOM generally increased with age.
- The proportion of children and young people with no ear conditions generally increased with age.



## Changes among children and young people who received 2 or more services between 2012 and 2019


This section presents information about changes in ear conditions, hearing loss and hearing impairment among children and young people who received more than 1 audiology service during the hearing health outreach services treatment pathway. This is one way to understand the effectiveness of the hearing health outreach services in terms of improving outcomes. It is important to keep in mind that changes may also be partially attributed to the natural progression of the disease as children and young people grow older. To measure changes over time and observe the outcomes of treatment, this analysis includes only children and young people who received 2 or more services with a minimum interval of 3 months between their first and last service. A total of 3,965 children and young people met this criteria.

For almost all conditions, there was an improvement over time. Figure 3.3 shows the change in proportion of ear conditions between first and last services:

- The number of children and young people with at least 1 type of ear condition decreased from 3,005 at their first service to 2,039 at their last service.
- The proportion of children and young people with at least 1 type of ear condition decreased by 25 percentage points between the first and the last service (from 76% to 51%).
- There was a decrease in the proportion of children with an ear condition between first and last services for the majority of middle ear conditions, except for CSOM without discharge, for which results were similar to those in 2018.



**Figure 3.3: Change in proportion of ear conditions between first and last hearing health service, among children who received at least 2 services July 2012 and December 2019**



(a) Otitis media includes OME, CSOM with discharge, CSOM without discharge and AOM. The sum of these 4 conditions may exceed the total of otitis media as service recipients may have more than 1 ear condition.

(b) Other ear conditions includes grommets, reduced ear drum movement or retracted ear drum, and other ear conditions.

Source: table S3.3.

## Ear conditions among CNS service recipients

Children and young people who receive CNS services have a prioritised need for care. As such, it is important to examine these children and young people separately to track their hearing health as they move through the program.

In 2019, among the 1,008 children who received a CNS service and consented to sharing information with the AIHW, 688 children and young people (68%) were diagnosed with at least 1 ear condition at their first CNS visit. Thirty percent were found to have no ear condition (information was missing for the remaining 2%).

The most commonly diagnosed conditions were:

- OME (30%)
- Eustachian tube dysfunction (19%)
- AOM with discharge (8%).

Compared with 2018, more children had OME and AOM, but fewer children had ETD and CSOM with discharge. This indicated that more children in 2019 had been treated at the acute phase of otitis media than in 2018.


**Find out more in Table S3.5.**

# Hearing status

Two measures of hearing status are used in this report: hearing loss and hearing impairment. 'Hearing loss' may affect 1 ear (unilateral) or both ears (bilateral). 'Hearing impairment' is based on the ear with the better hearing, meaning that children and young people with unilateral hearing loss are not defined as having a hearing impairment. Only those with bilateral hearing loss are classified according to the degree of hearing impairment.

Figure 3.4 shows the number and proportion of children and young people with hearing loss and hearing impairment among service recipients in 2019, and the relationship between the two.

**Figure 3.4: Hearing loss and impairment among children and young people who received audiology outreach services (including CNS service recipients), 2019**



Sources: tables S2.1, S3.6, S3.8.

## Hearing loss

There are 3 types of hearing loss: conductive, sensorineural and mixed (Box 3.2). Among the 1,896 children and young people who received audiology outreach services in 2019, 42% had hearing loss:

- 27% had conductive hearing loss
- 0.6% had sensorineural hearing loss
- 0.7% had mixed hearing loss
- 14% could not be determined.

About 1,170 children and young people who received outreach audiology were visited by CNSs in 2019. Among these children, 48% had some form of hearing loss—29% bilateral and 19% unilateral.


Among children who received CNS services between July 2012 and December 2019, there was a remarkable reduction—31 percentage points—in the proportion of children with hearing loss.

*Find out more in Table S3.7 and Table S3.9.*

## Variation by age and sex

Among children and young people who received an audiology service in 2019, the proportion with hearing loss differed between age groups (Figure 3.5). Among young people aged 3–5, 47% had hearing loss, the highest proportion of any age group.

**Figure 3.5: Proportion of children and young people with hearing loss who received audiology outreach services in 2019, by age group**



Source: table S3.10.

### Box 3.2: Types of hearing loss

#### Conductive hearing loss:

A deviation of hearing threshold from normal range associated with reduced conduction of sound through the outer ear, tympanic membrane (eardrum) or middle ear, including ossicles (middle ear bones).


#### Sensorineural hearing loss:

A deviation of hearing threshold from the normal range, attributable to problems in the inner ear or vestibulocochlear nerve.

**Mixed hearing loss:** Hearing loss that has conductive and sensorineural components combined.

The proportion of hearing loss was the same among girls and boys in 2019 (Figure 3.6).

**Figure 3.6: Hearing loss among children and young people in 2019, by sex**




### Hearing loss among children and young people with ear conditions

Proportions of hearing loss were much higher among children and young people who had ear conditions than among those with no ear conditions (Figure 3.7).

**Figure 3.7: Hearing loss among children and young people in 2019 with ear conditions**

In 2019, among children who were diagnosed with **at least 1** ear condition:




Source: Table S3.11.

### Trends

Among children and young people who received an audiology service, the proportion with hearing loss decreased from 55% in July–December 2012 to 42% in 2019 (Figure 3.8).

**Figure 3.8: Hearing loss among service recipients over time**

When comparing service recipients from July–December 2012 and 2019:




Note: About 11% of service recipients whose hearing loss information were 'missing' from the data. 'Missing' includes not tested, not stated, unsure or invalid responses.

Source: Table S3.6.

## Hearing impairment

In this report, hearing impairment is defined as the degree of impairment associated with hearing loss in the 'better hearing ear', using a scale of mild, moderate, severe and profound (Australian Hearing, cited in Access Economics 2006). In 2019, 445 (24%) children and young people who received an audiology service had a hearing impairment (Figure 3.9).

**Figure 3.9: Hearing impairment among service recipients in 2019**




Source: Table S3.8.

## Variation by age and sex


The proportion of boys with hearing impairment was slightly higher than the proportion of girls with hearing impairment in 2019 (Figure 3.10).

**Figure 3.10: Hearing impairment in 2019, by sex**



In 2019, the proportion of children and young people with no hearing impairment generally rose with age, and hearing impairment tended to be more severe in younger than in older age groups (3.11).

**Figure 3.11: Hearing impairment in 2019 for younger ages**




Source: Table S3.13.

The shifts with age in the severity of hearing impairment can be partly explained by the effects of both medical intervention and natural development (whereby children typically grow out of ear conditions and associated hearing loss). At older ages (3–5 onwards), it is likely that medical treatment and interventions for those with ear conditions help reduce the severity of hearing impairment, and therefore reduce the proportion of children and young people with impairment.

## Trends

Among children and young people who received outreach audiology services between July 2012 and December 2019, the proportion with a hearing impairment decreased from 40% to 24% (Figure 3.12).

**Figure 3.12: Degree of hearing impairment among children and young people, July 2012 to December 2019**



Source: Table S3.8.

Some of the hearing impairment decrease over time could be due to the natural history of ear disease, as previously discussed. However, it is more likely that the observed decrease is attributable to the increasing effectiveness of hearing health services and medical interventions. It is difficult to attribute such a large decrease in hearing impairment over a short period of time solely to the natural progression of the disease. Overall, the effectiveness of these services can be measured only through an evaluation program, which is beyond the scope of this report.


## Changes among children who received 2 or more services between 2012 and 2019

One way to examine hearing health changes is by matching first and last visits among children and young people as they move through the health system. Individual children who received multiple audiology services were tracked to determine whether their hearing health changed (see Box 3.3).

Among the 3,480 children and young people who received 2 or more outreach audiology services between July 2012 and December 2019, hearing health generally improved over the period (Figure 3.13):

- 1,553 children (61%) had an improved hearing loss status and the status of 105 (4%) had deteriorated.
- Among the 1,647 children with hearing impairment, 1,165 (71%) had improved capability and for 81 (5%) capability had deteriorated.

**Figure 3.13: Change in hearing loss and impairment among children who received at least 2 audiology services between July 2012 and December 2019**



Sources: tables S3.14, S3.15.

### Box 3.3: Change in hearing capability across services

#### Hearing loss


**Improved**—a change in hearing loss status: (1) from bilateral hearing loss to unilateral hearing loss or no hearing loss and (2) from unilateral hearing loss to no hearing loss.

**Deteriorated**—a change in hearing loss status: (1) from no hearing loss to unilateral or bilateral hearing loss or (2) from unilateral hearing loss to bilateral hearing loss.

#### Hearing impairment

**Improved**—movement to a lower degree of hearing impairment (for example, from profound hearing impairment to severe, moderate or mild hearing impairment).

**Deteriorated**—movement to a higher degree of hearing impairment (for example, from mild hearing impairment to moderate, severe or profound).



A number of factors might contribute to the observed improvements in hearing health. These include:

- the effectiveness of medical interventions
- the effect of health promotion activities in:
  - increasing awareness and knowledge of hearing health among families
  - improving the acceptance of and attendance at audiology services provided by outreach teams
- the natural improvement in the condition as children and youth get older
- the confounding factor of ‘missing’ and ‘not tested’ records, and missing information from children and youth for whom consent was not obtained.





# 4

## **Demand for ear and hearing health services and other follow-up services**

The Australian Government-funded activities in the Northern Territory public hearing health system are valuable for children and young people in the Northern Territory. Since 2012, the number of children and young people accessing services has increased, and consent rates have increased as well. Improvements in hearing health status across the years have also been consistently demonstrated. However, there is still a high demand for audiology and ENT services.


In addition to these hearing health services, children can be referred to other follow-up services. Follow-up can encompass a wide range of services, from having a hearing aid fitted by Hearing Australia, to medical treatment such as ear cleaning or the need for ENT surgery.

## Follow-up services required after audiologist visits

In 2019, 67% of children and young people required at least 1 further action following their audiology visit. (Percentages in the infographic Figure 4.1 below sum to over 67% because some children required more than 1 referral).

**Figure 4.1: Follow-up services required after audiology visits**

In 2019, 1,896 children and young people received an audiology service. Among these:



Source: Table S4.1.

## Follow-up services required after ENT teleotology services

In 2019, 464 children and young people (65%) were given a recommendation for at least 1 further action following an ENT teleotology service. Two types of action are recommended by ENT specialists:

- surgery—the most common types of surgery recommended were myringoplasty and myringotomy (see Glossary).
- further follow-up—this is the main ENT action recommended through the ENT teleotology service. This was primarily for an ENT review or an audiological assessment.




## Waiting list for audiology services and ENT teleotology services

Although a number of hearing health services have been provided, the demand for audiology services and ENT teleotology service remains high among Indigenous children and young people in the Northern Territory for the following reasons:

- ear disease is a chronic condition that often requires multiple service events over a number of years
- regular training and education sessions on ear disease and hearing loss are being delivered in the community, which has resulted in increased knowledge and awareness of ear disease and the impacts of hearing loss
- more effective identification of ear disease in the community and through Hearing Services auditing processes results in an increase in referrals
- restricted capacity of Hearing Services to provide frequent service to some communities due to availability of ear and hearing health-care staff
- cultural events, inclement weather, and school and public holidays restrict the capacity of the teams to travel to remote communities all year round
- the geographical location and vast spread of Indigenous communities make access to services difficult
- the high rates of mobility between communities for families can hinder efforts to locate children in a timely manner and provide appropriate ear and hearing health.

At the end of 2019, there were 3,459 children and young people on the audiology waiting list. Of these, 1,726 were new referrals for audiology services and 1,733 were recalls for further audiology follow-up.

In addition, 2,458 children and young people were waiting for ENT teleotology services in the Northern Territory. Of these, 1,025 were new referrals either from Primary Health Care physicians or from the audiology pathway, and required ENT intervention. There were 1,433 children and young people who had been reviewed but required additional follow-up by ENT teleotology services.



# 5

## Regional analysis




The Northern Territory has 2 main health service delivery regions: the Top End Health Service (TEHS) and the Central Australia Health Service (CAHS). This section presents the hearing health status of the children and young people who received outreach audiology, CNS or ENT teleotology services in these 2 regions. In 2019, 1,356 children and young people within the TEHS and 500 within the CAHS received an audiology service.

## Hearing loss

Patterns over the years were generally similar for children in the TEHS and the CAHS between July 2012 and December 2016. However, in 2017 the difference in the proportion of children with hearing loss grew between the regions. In 2019, the gap between the TEHS and the CAHS narrowed again (Figure 5.1):

- Children in the CAHS had a higher proportion of hearing loss (44%, or 220 children) than those in the TEHS (41%, or 557 children).
- The proportion of children and young people with no hearing loss was higher in the TEHS (47%, or 635 children) than in the CAHS (45%, or 225 children).


**Figure 5.1: Hearing loss among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS) July 2012 to December 2019**



Source: Table S5.1.

When looking at specific types of hearing loss in 2019 (Figure 5.2), the proportions between children in the TEHS and the CAHS are generally similar.

Figure 5.2: Types of hearing loss among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS) July 2012 to December 2019



Sources: tables S5.1, S5.2.


## Hearing impairment

Overall, for both regions from July 2012 to December 2019:

- the proportion of children and young people with hearing impairment decreased over time, (Figure 5.3) by:
  - 15 percentage points for the TEHS
  - 19 percentage points for the CAHS
- the proportion of children and young people with no hearing impairment increased over the years by:
  - 16 percentage points for the TEHS
  - 10 percentage points for the CAHS.

From July 2012 to December 2016, hearing impairment patterns across the years were similar for children and young people who received services in the TEHS and the CAHS, but there was a deviation from these patterns from 2017. The gap between the TEHS and CAHS then narrowed in 2019, and children and young people in the TEHS had similar proportions of hearing impairment (24% of the CAHS). Proportions of children and young people with mild hearing impairment were higher in the CAHS than in the TEHS (18% compared with 16%) (Figure 5.4). The proportions of children with moderate hearing impairment were similar in the 2 regions.

**Figure 5.3: Hearing impairment among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS) July 2012 to December 2019**



Sources: Table S5.3.



Figure 5.4: Types of hearing impairment among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS), July 2012 to December 2019




Source: Table S5.3.

## Ear conditions

Between July 2012 and December 2019, the proportion of children and young people who received an outreach audiology, CNS or ENT teleotology service and who had an ear condition decreased overall in both regions—from 65% to 61% in the TEHS and from 73% to 66% in the CAHS (Figure 5.5).

**Figure 5.5: Presence of ear conditions among children and young people who received outreach audiology, CNS or ENT teleotology services in the Top End (TEHS) and Central Australia (CAHS) July 2012 to December 2019**




Source: Table S5.4.


The distribution of types of ear condition varied over time in both regions (Figure 5.6). Notably, from July 2012 to December 2019:

- the proportion of children with OME increased in the TEHS, from 21% to 25%, but decreased from 36% to 24% in the CAHS
- the proportion of children with CSOM with discharge decreased in the TEHS, from 17% to 9%, but increased in the CAHS from 8% to 10%
- the proportion of children with ETD increased slightly from 16% to 17% for the TEHS and decreased slightly from 18% to 17% for the CAHS.

Figure 5.6: Distribution of specific ear conditions among children and young people who received a service in Central Australia (CAHS) and the Top End (TEHS) July 2012 to December 2019



Source: Table S5.4.



# 6

## Progress against benchmarks

The Hearing Health Program uses performance indicators and benchmarks to monitor outcomes. Targets are set jointly by the Australian and Northern Territory departments of health through the Northern Territory Health Implementation Plan (Council on Federal Financial Relations 2016).

Some targets are annual, and others are set for a longer period. Most of the targets for 2019 were met or exceeded. Some targets are difficult to measure at this point as they are designed to be met within 3 years.


## Service delivery

The annual benchmark for hearing health service delivery is measured by the number of services per year. From 2012 to 2019, service delivery targets for audiology services, CNS services and hearing health promotion services were either met or exceeded. Note that the Northern Territory Health Implementation Plan does not include targets for ENT services.

**Indicator:**  
**Audiology services provided**

The number of **audiology services** per year


**Figure 6.1: Number of audiology services provided, 2013-2019**



**Indicator:**  
**CNS services provided**

The number of children receiving complex case management services from **CNSs** working with primary health-care services


**Figure 6.2: Number of children receiving CNS services, 2013-2019**



**Indicator:**  
**Health promotion**

Delivery of hearing health promotion or training services and activities

**Figure 6.3: Number of activities/sessions provided in 2019**




## Health outcomes—hearing impairment

Health outcomes in this program are measured every 3 years (July 2012–June 2015, July 2015–December 2018 and January 2019–June 2021). At the beginning of each period, the benchmarks for health outcomes were raised. All benchmarks in the first 2 periods have been met. As 2019 is the first year of the new period, the information on this page is based on the data from 1 year only and does not indicate whether the benchmarks have been met.

### Indicator: Hearing impairment

Proportion of children tested between January 2019 and December 2019 with moderate or severe conductive hearing impairment

**Figure 6.4: Proportion of children with moderate/severe conductive hearing impairment, January 2019 to December 2019**




## Health outcomes—middle ear conditions


### Indicator: Children with CSOM

Proportion of children who received an audiology check or CNS service between January 2019 and December 2019 who were found to have CSOM

**Figure 6.5a: Proportion of children (aged 0–5) with CSOM, January 2019 to December 2019**




**Figure 6.5b: Proportion of children (aged 6–15) with CSOM, January 2019 to December 2019**




### Indicator: Children with dry perforation

Proportion of children who received an audiology check or CNS service between January 2019 and December 2019 who were found to have dry perforation

**Figure 6.6a: Proportion of children (aged 0–5) with dry perforation, January 2019 to December 2019**



**Figure 6.6b: Proportion of children (aged 6–15) with dry perforation, January 2019 to December 2019**





# Appendix A: About the Hearing Health Program data collections

## Data collection, management and reporting

The AIHW was commissioned by the Department of Health to collect, manage and report on data from ear and hearing health outreach services in the Northern Territory.

These data are collected using paper data collection forms. Health professionals responsible for providing services complete a form with information about the child or young person's demographic characteristics; types of services provided; community where the service was provided; date of service; examination results; and medical interventions and recommendations.

How much data the AIHW receives on each child or young person depends on whether the child or young person's parent or guardian provides consent to share the information. There are 2 scenarios for the provision of data under the consent requirements:

- If consent is given, all de-identified data are sent to the AIHW.
- If consent is not given, a limited amount of aggregate information is provided to the AIHW. This includes the number of services provided and the number of children and young people receiving a service by 5-year age group, sex, and community where the service was provided.

Throughout this report, the term 'services' refers to occasions of service. A child or young person may receive a number of services and have more than 1 record in each data collection. Each record in the collection corresponds to a single service, not to a single person.

More information on each of the Hearing Health Program data collections can be found in the Reference Material online at <https://www.aihw.gov.au/reports/indigenous-australians/hearing-health-outreach-services/formats>.



## Appendix B: Data quality statement

For all Hearing Health Program data collections, the population included is not a random sample, nor is it representative of all Aboriginal and Torres Strait Islander children and young people in the Northern Territory. The outreach audiology and ENT teleotology services are available to all Indigenous children and young people, but not all of them access these services. The CNS program is available only to Indigenous children who have a referral from a health professional.

As well, some of these services are more commonly accessed by individuals in remote areas. Therefore, results of analyses cannot be generalised to all Indigenous children and young people in the Northern Territory.

Due to differences in the scope of the programs covered in previous AIHW hearing health reports, analyses from individual reports should not be compared with analyses in subsequent annual reports.

### Outreach audiology data collection summary

- This data collection included about 8,000 children and young people, aged under 21, who received Northern Territory outreach audiology services between July 2012 and December 2019. This accounted for about 26% of the Northern Territory's Indigenous population of this age group (but was not a random sample).
- Hearing loss status was missing for about 12% of service participants who completed audiology assessments in 2019, and this should be considered when using and interpreting hearing health data.
- The full data quality statement for the audiology data collection can be found online at: <https://meteor.aihw.gov.au/content/index.phtml/itemId/719804>.

### ENT teleotology data collection summary

- This data collection included over 3,600 children and young people who were aged under 21 and received ENT teleotology services between July 2012 and December 2019. This accounted for about 11% of the Northern Territory Indigenous population of this age group (but was not a random sample).
- The methods of assessment used at ENT teleotology services differ from those for face-to-face consultations. Results of tests and subsequent diagnoses from teleotology services may be affected by the method of service delivery.
- The full data quality statement for ENT teleotology data collection can be found online at: <https://meteor.aihw.gov.au/content/index.phtml/itemId/719809>.

## CNS data collection summary

- The data collection includes over 4,500 children aged under 21 who received CNS services between July 2012 to December 2019. This accounted for about 13% of the Northern Territory's Indigenous population of this age group (but was not a random sample).
- Prior to 2016, rates of non-consent were high for the CNS program (20% of services and 21% of children in 2015). However, there have been improvements in non-consent rates over time, and in 2018 there were no cases of non-consent. This should be considered when interpreting CNS program analyses.
- The full data quality statement for the CNS data collection can be found online at: <https://meteor.aihw.gov.au/content/index.phtml/itemId/719806>.

## Child Health Check Initiative (CHCI) data collections summary

- The data from August 2007 to July 2012 included in this report is from the CHCI data collections.
- Children who received child health checks or follow-up services were not a random sample of Indigenous children in the Northern Territory. Health checks and services were available only to children in prescribed areas of the Northern Territory and were provided on a voluntary basis.
- The full data quality statement for the CHCI data collections can be found online at: <https://meteor.aihw.gov.au/content/index.phtml/itemId/480005>.



## Acknowledgments

This report was prepared by Ms Bin Tong, Ms Ruth Penm and Dr Bina Gubhaju of the Australian Institute of Health and Welfare. The authors acknowledge Mr Geoff Callaghan and Dr Fadwa Al-Yaman of the AIHW for their comments and guidance throughout the preparation of this report.

The Department of Health provided funds to support data collection and report production, and members of the Indigenous Health Division are gratefully acknowledged for their comments on this report.

Special thanks go to the many clinicians and administrators who supported the hearing health data collections included in this report. The assistance and cooperation of the Northern Territory Department of Health is greatly appreciated.

The authors also thank Professor Harvey Coates of the School of Paediatrics and Child Health at the University of Western Australia and Dr Kelvin Kong of John Hunter Children's Hospital for reviewing and commenting on the report.

# Abbreviations

ABS	Australian Bureau of Statistics
AIHW	Australian Institute of Health and Welfare
AOM	acute otitis media
CAHS	Central Australia Health Service
CHCI	Child Health Check Initiative
CNS	Clinical Nurse Specialist
CSOM	chronic suppurative otitis media
CtG	Closing the Gap program
ENT	ear, nose and throat
ETD	Eustachian tube dysfunction
HP	Health priority
NT	Northern Territory
NTER	Northern Territory Emergency Response
NTRAI	Northern Territory Remote Aboriginal Investment
OME	otitis media with effusion
SFNT	National Partnership Agreement on Stronger Futures in the Northern Territory
TEHS	Top End Health Service
WHO	World Health Organization

# Symbols

$\geq$	greater than or equal to
$\leq$	less than or equal to
$<$	less than
$>$	greater than

# Glossary

**acute otitis media:** The general term for both acute otitis media without perforation and acute otitis media with perforation. It is the presence of fluid behind the eardrum plus at least 1 of the following: bulging eardrum, red eardrum, recent discharge of pus, fever, ear pain or irritability. A bulging eardrum, recent discharge of pus, and ear pain are the most reliable indicators of acute otitis media.

**aural toilet:** A procedure where an ear, nose and throat surgeon clears wax, debris or foreign bodies from the ear canal. It is often used in treating patients with recurrent infections of the ear canal.

**bilateral hearing loss:** Hearing loss in both ears.

**chronic suppurative otitis media (CSOM) with discharge:** A persistent **suppurative** discharge from the middle ear through a tympanic membrane (ear drum) perforation for more than 6 weeks. Importantly, the diagnosis of CSOM with discharge is appropriate only if the tympanic membrane perforation is seen and if it is large enough to allow the discharge to flow out of the middle ear space.

**chronic suppurative otitis media without discharge:** The presence of a perforation (hole) in the eardrum without evidence of discharge or fluid behind the eardrum. It is also known as 'inactive chronic suppurative otitis media', and also as 'dry perforation'.

**conductive hearing loss:** A deviation of hearing threshold from the normal range associated with reduced conduction of sound through the outer ear, tympanic membrane (eardrum) or middle ear, including ossicles (middle ear bones).

**Eustachian tube dysfunction:** Negative middle ear pressure associated with compromised equalisation impeding middle ear function and sometimes causing middle ear fluid accumulation.

**grommet:** A small tube surgically placed across the eardrum to re-establish ventilation to the middle ear. It is also called 'ventilation tube', 'pressure equalisation (PE) tube', or a 'tympanostomy tube'.

**hearing:** The sense for perceiving sounds; includes regions within the brain where the signals are received and interpreted.

**hearing impairment:** Describes the degree of impairment associated with hearing loss in the 'better hearing ear', using a scale of mild, moderate, severe and profound. It is based on degree of deviation from normal thresholds in the 'better ear', calculated as a 3-frequency average of the threshold of hearing (in dB HL)—500 Hz, 1000 Hz and 2000 Hz.

**hearing loss:** Any hearing threshold response (using audiometry) outside the normal range, at any sound stimuli, in either ear. Hearing loss in a population describes the number of people who have abnormal hearing. Hearing loss may affect 1 ear (unilateral) or both ears (bilateral).

**mild hearing impairment:** On average, the quietest sounds that people can hear with their 'better' ear are 16–30 dB HL in soundproof conditions and 26–35 dB HL in non-soundproof conditions. These people are able to hear and repeat words spoken in a normal voice at 1 metre. Counselling and hearing aids may be needed.

**mixed hearing loss:** Hearing loss that has **conductive** and **sensorineural** components combined.

**moderate hearing impairment:** On average, the quietest sounds that people can hear with their 'better' ear are 31–60 dB HL in soundproof conditions and 36–60 dB HL in non-soundproof conditions. These people are able to hear and repeat words spoken in raised voice at 1 metre and have difficulty keeping up with conversations without using a hearing aid.

**myringoplasty:** The repair of a perforation of the tympanic membrane (eardrum).

**myringotomy:** Surgical incision into the eardrum, to relieve pressure or drain fluid.

**otitis media:** All forms of inflammation and infection of the middle ear. Active inflammation or infection is nearly always associated with a middle ear effusion (fluid in the middle ear space).

**otitis media with effusion (OME):** The presence of an intact eardrum and middle ear fluid without symptoms or signs of acute infection. Other terms used to describe OME include 'glue ear', 'serous otitis media' and 'secretory otitis media'. OME may be episodic or persistent.

**profound hearing impairment:** On average, the quietest sounds that people can hear with their better ear are 91+ dB HL either in soundproof conditions or non-soundproof conditions. These people are unable to hear and understand even a shouted voice. Hearing aids may help in understanding words. Additional rehabilitation is needed, and cochlear implants, lip-reading and sometimes signing are necessary.

**sensorineural hearing loss:** A deviation of hearing threshold from the normal range, attributable to problems in the inner ear or vestibulocochlear nerve.

**severe hearing impairment:** On average, the quietest sounds that people can hear with their better ear are 61–90 dB HL, either in soundproof conditions or non-soundproof conditions. These people are able to hear some words when shouted into the 'better' ear. Hearing aids are needed; if no hearing aids are available, lip-reading and signing may be necessary.

**suppurative:** Pus produced in response to inflammatory bacterial infections.

**teleotology:** Method of offsite service delivery whereby specialists and audiologists provide full diagnostic hearing assessments, assess middle ear function, diagnose middle ear conditions and recommend further actions and treatment based on information provided to them electronically by an audiologist or an ENT nurse consultant.

**unilateral hearing loss:** Hearing loss in 1 ear.

## References

- ABS (Australian Bureau of Statistics) 2016. National Aboriginal and Torres Strait Islander Social Survey, 2014–15. ABS cat. no. 4714.0. Canberra: ABS. Viewed 10 June 2016, <<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4714.02014-15>>.
- Access Economics 2006. Listen hear: the economic impact and cost of hearing loss in Australia. Sydney: Access Economics.
- AIHW (Australian Institute of Health and Welfare) 2012. Northern Territory Emergency Response Child Health Check Initiative—follow-up services for oral and ear health: final report, 2007–2012. Cat. no. DEN 223. Canberra: AIHW.
- AIHW 2014a. Access to primary health care relative to need for Indigenous Australians. Cat. no. IHW 128. Canberra: AIHW.
- AIHW 2014b. Australia's health 2014. Australia's health series no. 14. Cat. no. AUS 178. Canberra: AIHW.
- AIHW 2014c. Stronger Futures in the Northern Territory: hearing health services 2012–2013. Cat. no. IHW 117. Canberra: AIHW.
- AIHW 2017a. Aboriginal and Torres Strait Islander health performance framework 2017: supplementary online tables. Cat. no. WEB 170. Canberra: AIHW.
- AIHW 2017b. Northern Territory Outreach Hearing Health Program: July 2012 to December 2016. Cat. no. IHW 189. Canberra: AIHW.
- AIHW & DoHA (Department of Health and Ageing) 2009. Progress of the Northern Territory Emergency Response Child Health Check Initiative: update on results from the Child Health Check and follow-up data collections. Cat. no. IHW 28. Canberra: AIHW.
- Council on Federal Financial Relations 2016. Northern Territory Health Implementation Plan: National Partnership on Northern Territory Remote Aboriginal Investment. Canberra: Council on Federal Financial Relations.
- Jervis-Bardy J, Sanchez L & Carney AS 2014. Otitis media in Indigenous Australian children: review of epidemiology and risk factors. *Journal of Laryngology and Otology* 128(S1):S16–27.
- Moeller MP 2000. Early intervention and language development in children who are deaf and hard of hearing. *Pediatrics* 106(3):e43.
- WHO (World Health Organization) 1996. Prevention of hearing impairment from chronic otitis media: report of a WHO/CIBA Foundation workshop, London, United Kingdom, 19–21 November 1996. Geneva: World Health Organization.



## List of tables

Table 1: Progress against benchmarks, 2019 .....	vii
--	-----

## List of boxes

Box 2.1: Hearing health priority (HP) categories.....	13
Box 3.1: Types of ear conditions.....	15
Box 3.2: Types of hearing loss .....	21
Box 3.3: Change in hearing capability across services.....	25

## List of figures

Figure 1.1: Clinical path of children and young people moving through the Northern Territory public hearing health system .....	4
Figure 2.1: Ear health education, promotion and prevention training sessions delivered in 2019.....	7
Figure 2.2: Number of outreach audiology services and service recipients, July 2012–December 2019 .....	8
Figure 2.3: Outreach audiology service recipients in 2019, by sex.....	9
Figure 2.4: Outreach audiology service recipients in 2019, by age group .....	9
Figure 2.5: Number of ENT teleotology services and service recipients, July 2012–December 2019 .....	10
Figure 2.6: ENT teleotology service recipients in 2019, by sex .....	11
Figure 2.7: ENT teleotology service recipients in 2019, by age group.....	11
Figure 2.8: Number of CNS services and service recipients, July 2012–December 2019 .....	12
Figure 2.9: Clinical Nurse Specialist service recipients in 2019, by sex.....	13
Figure 2.10: Hearing health priority (closely linked to age) among CNS service recipients, 2019 .....	13
Figure 3.1: Types of ear conditions among children and young people who received an audiology, CNS or ENT teleotology service in 2019.....	16
Figure 3.2: Most common ear conditions in children and young people who received at least 1 audiology outreach, CNS or ENT teleotology service in 2019, by age group .....	17
Figure 3.3: Change in proportion of ear conditions between first and last hearing health service, among children who received at least 2 services between July 2012 and December 2019.....	18




Figure 3.4: Hearing loss and impairment among children and young people who received audiology outreach services (including CNS service recipients), 2019.....20

Figure 3.5: Proportion of children and young people with hearing loss who received audiology outreach services in 2019, by age group.....21

Figure 3.6: Hearing loss among children and young people in 2019, by sex.....22

Figure 3.7: Hearing loss among children and young people in 2019 with ear conditions.....22

Figure 3.8: Hearing loss among service recipients over time.....22

Figure 3.9: Hearing impairment among service recipients in 2019.....23

Figure 3.10: Hearing impairment in 2019, by sex.....23

Figure 3.11: Hearing impairment in 2019 for younger ages.....23

Figure 3.12: Degree of hearing impairment among children and young people, July 2012 to December 2019.....24

Figure 3.13: Change in hearing loss and impairment among children who received at least 2 audiology services between July 2012 and December 2019.....25

Figure 4.1: Follow-up services required after audiology visits.....28

Figure 5.1: Hearing loss among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS) July 2012–2019.....31

Figure 5.2: Types of hearing loss among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS).....32

Figure 5.3: Hearing impairment among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS).....33

Figure 5.4: Types of hearing impairment among children and young people who received outreach audiology services in Central Australia (CAHS) and the Top End (TEHS), July 2012–2019.....34

Figure 5.5: Presence of ear conditions among children and young people who received outreach audiology, CNS or ENT teleotology services in the Top End (TEHS) and Central Australia (CAHS).....35

Figure 5.6: Distribution of specific ear conditions among children and young people who received a service in Central Australia (CAHS) and the Top End (TEHS).....36



Figure 6.1: Number of audiology services provided, 2013–2019.....	38
Figure 6.2: Number of children receiving CNS services, 2013–2019.....	38
Figure 6.3: Number of activities/sessions provided in 2019.....	38
Figure 6.4: Proportion of children with moderate/severe conductive hearing impairment, January 2019 to December 2019.....	39
Figure 6.5a: Number of children (aged 0–5) with CSOM, January 2019 to December 2019 .....	39
Figure 6.5b: Number of children (aged 6–15) with CSOM, January 2019 to December 2019 .....	39
Figure 6.6a: Number of children (aged 0–5) with dry perforation, January 2019 to December 2019.....	39
Figure 6.6b: Number of children (aged 6–15) with dry perforation, January 2019 to December 2019 .....	39


## Related publications

The following AIHW publications on the Child Health Check Initiative Closing the Gap; National Partnership on Stronger Futures in the Northern Territory; and Northern Territory Remote Aboriginal Investment hearing health programs may be of interest:

- AIHW (Australian Institute of Health and Welfare) 2010. Health and wellbeing of young Australians: indicator framework and key national indicators. Bulletin no. 77. Cat. no. AUS 123. Canberra: AIHW.
- AIHW 2011. Ear and hearing health of Aboriginal and Torres Strait Islander children in the Northern Territory. Cat. no. IHW 60. Canberra: AIHW.
- AIHW 2012. Northern Territory Emergency Response Child Health Check Initiative: follow-up services for oral and ear health, final report, August 2007–June 2012. Cat. no. DEN 223. Canberra: AIHW.
- AIHW 2014. Stronger Futures in the Northern Territory: hearing health services 2012–2013. Cat. no. IHW 117. Canberra: AIHW.
- AIHW 2015. Hearing health outreach services to Aboriginal and Torres Strait Islander children and young people in the Northern Territory: 2012–13 to 2014–15. Cat. no. IHW 163. Canberra: AIHW.
- AIHW 2015. Hearing health outreach services to Indigenous children and young people in the Northern Territory: 2012–13 and 2013–14. Cat. no. IHW 149. Canberra: AIHW.
- AIHW 2017. Northern Territory Outreach Hearing Health Program: July 2012 to December 2016. Cat. no. IHW 189. Canberra: AIHW.
- AIHW 2017. Northern Territory Remote Aboriginal Investment: Ear and Hearing Health Program—July 2012 to June 2016. Cat. no. IHW 176. Canberra: AIHW.
- AIHW 2018. Northern Territory Outreach Hearing Health Program: July 2012 to December 2017. Cat. no. IHW 203. Canberra: AIHW.
- AIHW 2019. Hearing health outreach services for Aboriginal and Torres Strait Islander children in the Northern Territory: July 2012 to December 2018. Cat. No. IHW 213. Canberra: AIHW.

These reports can be downloaded free from the AIHW website at <http://www.aihw.gov.au/publications>.

The website also provides information on ordering printed copies.



This report presents information on hearing health outreach services provided to Aboriginal and Torres Strait Islander children and young people in the Northern Territory. It shows that in 2019, 2,156 audiology; 770 ear, nose and throat teleotology; and 1,119 Clinical Nurse Specialist services were provided. Among children and young people who received treatment, 61% had improved hearing loss and 71% had improved hearing impairment.

[aihw.gov.au](http://aihw.gov.au)



Stronger evidence,  
better decisions,  
improved health and welfare

