



Australian Government
Australian Institute of
Health and Welfare

AIHW

National Core Maternity Indicators 2017

Summary report



National Core Maternity Indicators 2017

Summary report



The Australian Institute of Health and Welfare is a major national agency whose purpose is to create authoritative and accessible information and statistics that inform decisions and improve the health and welfare of all Australians.

© Australian Institute of Health and Welfare 2019 

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC BY 3.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at <www.aihw.gov.au/copyright/>. The full terms and conditions of this licence are available at <<http://creativecommons.org/licenses/by/3.0/au/>>.

ISBN 978-1-76054-609-0 (Online)

ISBN 978-1-76054-610-6 (Print)

Suggested citation

Australian Institute of Health and Welfare 2019. National Core Maternity Indicators 2017: summary report. Cat. no. PER 104. Canberra: AIHW.

Australian Institute of Health and Welfare

Board Chair

Mrs Louise Markus

Chief Executive Officer

Mr Barry Sandison

Any enquiries about or comments on this publication should be directed to:

Australian Institute of Health and Welfare

GPO Box 570

Canberra ACT 2601

Tel: (02) 6244 1000


Email: info@aihw.gov.au

Published by the Australian Institute of Health and Welfare.

Please note that there is the potential for minor revisions of data in this report.
Please check the online version at <www.aihw.gov.au> for any amendments.

Contents

Introduction.....	1
Antenatal period indicators.....	2
Tobacco smoking in pregnancy.....	2
Antenatal care in the first trimester.....	3
Labour and birth indicators.....	4
Birth method—selected women	4
Induction of labour for selected women giving birth for the first time	6
Women having their second birth vaginally whose first birth was by caesarean section.....	7
General anaesthetic for women giving birth by caesarean section	8
Episiotomy.....	9
Birth outcome indicators	10
Third and fourth degree tears	10
Apgar score of less than 7 at 5 minutes for births at or after term.....	11
Small babies among births at or after 40 weeks gestation	12
Data sources and methods	13
Acknowledgments.....	14
References.....	14



Introduction

In 2017, 301,095 women gave birth in Australia. Maternity services in Australia are provided by 8 state and territory health departments and a number of private providers. Each state and territory has differing care provision systems and care available to pregnant women and their babies.

National Core Maternity Indicators (NCMIs) provide information on measures of clinical activity and outcomes in relation to maternity care across Australia. The purpose of the indicators is to assist in improving the quality of maternity services in Australia by establishing baseline data for monitoring and evaluating practice change.

There are 12 indicators that cover data for the majority of women who gave birth in Australia in 2017 and are grouped into 3 broad topic areas:

- antenatal (during pregnancy) period,
- labour and birth, and
- birth outcomes.

Antenatal period indicators

Tobacco smoking in pregnancy

- a. in the first 20 weeks of pregnancy for all women giving birth
- b. after the first 20 weeks of pregnancy for all women who gave birth and reported smoking during pregnancy

Antenatal care in the first trimester for all women giving birth

Labour and birth indicators

Induction of labour for selected women giving birth for the first time

Caesarean section for selected women giving birth for the first time

Non-instrumental vaginal birth for selected women giving birth for the first time

Instrumental vaginal birth for selected women giving birth for the first time

Episiotomy for women having their first baby and giving birth vaginally

- a. without instruments to assist the birth
- b. assisted with instruments

General anaesthetic for women giving birth by caesarean section

Women having their second birth vaginally whose first birth was by caesarean section

Birth outcome indicators

Apgar score of less than 7 at 5 minutes for births at or after term

Small babies among births at or after 40 weeks gestation

Third and fourth degree tears

- a. for all vaginal first births
- b. for all vaginal births



For more information on each NCMI including an online interactive data display: see
[<https://www.aihw.gov.au/reports/mothers-babies/ncmi-data-visualisations/contents/summary>](https://www.aihw.gov.au/reports/mothers-babies/ncmi-data-visualisations/contents/summary)

Antenatal period indicators

Tobacco smoking in pregnancy


Smoking during pregnancy increases the risk of complications to both the baby and the mother. It increases the likelihood of premature birth, placental complications, stillbirth and neonatal death, as well as the risk of low birthweight (WHO 2013). Stopping smoking during pregnancy is key to reducing the risk of complications during pregnancy and birth as well as reducing adverse health outcomes for the baby.

This indicator examines the number of all women who gave birth that reported smoking in the first 20 weeks of pregnancy; and those that reported smoking after the first 20 weeks of pregnancy.

Key findings

- Fewer than 1 in 10 women who gave birth (28,573 or 9.6%) smoked in the first 20 weeks of pregnancy in 2017, a decrease from 12.9% in 2011.
- Around three-quarters (21,484 or 73.4%) of women who reported smoking in the first 20 weeks continued to smoke after 20 weeks of pregnancy.
- The proportion of women who smoked during the antenatal period was highest among those women living in the most disadvantaged areas.
- The proportion of women who stopped smoking after 20 weeks of pregnancy was lowest among those women living in the most disadvantaged areas.

Proportion of women who reported smoking during the antenatal period



Notes

1. Data excludes mothers whose smoking status was not stated.
2. Smoking during pregnancy is self-reported smoking of tobacco.
3. Due to differences in definitions and methods used for data collection, care must be taken in comparison across jurisdictions.

Find out more

To view data on smoking in pregnancy by state and territory of birth, mother's age, mother's Indigenous status, hospital sector, hospital size, remoteness of mother's usual residence and socioeconomic status, see the online interactive data display.

Antenatal care in the first trimester


Antenatal care is a planned visit between a pregnant woman and a midwife or doctor to assess and improve the wellbeing of the mother and baby throughout pregnancy. It does not include visits where the sole purpose is to confirm the pregnancy.

Antenatal care is associated with positive outcomes for the mother and child and increases the likelihood of receiving effective health interventions. It is particularly effective when started in the first trimester (before 14 weeks gestation), which often leads to better maternal health in pregnancy, fewer interventions in late pregnancy and positive health outcomes for the child (AIHW 2018; DoH 2018).

This indicator examines antenatal care visits in the first trimester for all women giving birth.

Key findings

- In 2017, 213,635 or 72.1% of women who gave birth received antenatal care in the first trimester, an increase from 187,755 or 65.9% in 2011.
- Younger women are less likely to receive antenatal care in the first trimester than older women. However, the gap between age groups has decreased steadily since 2012.



Labour and birth indicators

Birth method—selected women

In Australia, as well as around the world, there has been an increasing trend of intervention in the birthing process (caesarean section and instrumental vaginal birth), however, the appropriateness of such changes to clinical practice are debated (ACSQHC 2018). Women require clear information regarding the benefits of the intervention and the short- and long-term risks so that the woman is able make an informed decision as to whether such intervention is appropriate for her (NICE 2019).

There are three indicators related to birth method:

- Caesarean section for selected women giving birth for the first time.
- Non-instrumental vaginal birth for selected women giving birth for the first time.
- Instrumental vaginal birth for selected women giving birth for the first time.

Why 'selected women'?

Rather than the whole population, these indicators are based on 'selected women'. A range of factors and individual circumstances can influence the method of a baby's birth, including various health complications. Because of this variability, this data examines a group of women whose characteristics suggest they have a lower risk of such complications. This gives a better indication of what can be expected in 'standard' cases. Selected women are those that meet all of the following criteria:


- ✓ Were aged between 20–34 years
- ✓ Gave birth between 37–41 completed weeks of gestation
- ✓ Had a singleton baby who presented in the vertex (head down) position

These women made up about one-third of the approximate 301,000 women who gave birth in Australia in 2017.

For selected women giving birth for the first time:

- From 2004 to 2017, the rate of non-instrumental vaginal births decreased while instrumental vaginal births increased.
- The rate of caesarean sections increased from 25.3% in 2004 to 29.3% in 2017.

Birth method of selected women, 2004–2017




Notes

1. Data for selected women criteria, parity, were not available from Victoria for 2009.
2. Data for Tasmania from 2004 to 2012 were not available.

Non-instrumental vaginal birth	Instrumental vaginal birth	Caesarean section
Vaginal birth without intervention is seen by many as the preferred method of birth, as women tend to have fewer postnatal complications and are more physically able in the short term to care for their new babies (Rowland & Redshaw 2012).	The use of instruments (vacuum extraction cup or forceps) may be required to achieve a safe vaginal birth. Using instruments to assist birth is usually recommended when the condition of either the baby or the mother makes it less safe to allow time for unassisted birth to occur (RANZCOG 2016).	Caesarean section is one of the most common interventions in pregnancy and is safer now than in the past, however, a small risk of serious morbidity and mortality for both the mother and the baby remains (Betran et al. 2016; Villar et al. 2007; Keag et al. 2018).

Key findings

- Caesarean section was the most common birth method for selected women in private hospitals.
- 2 in 5 selected women giving birth in a private hospital had a caesarean section, compared to 1 in 4 selected women who gave birth in a public hospital.
- Those women living in the most disadvantaged socioeconomic areas were more likely to have a non-instrumental vaginal birth than those living in the least disadvantaged areas.
- In each of the five socioeconomic area quintiles, non-instrumental vaginal birth was the most common birth method for selected women, followed by caesarean section and instrumental vaginal birth.



Induction of labour for selected women giving birth for the first time


Induction is an intervention to stimulate the onset of labour. An induction of labour is undertaken when the risks of continuing the pregnancy are thought to be greater than the risks associated with being born (McDonnell 2011).

This indicator looks at induction of labour for selected women giving birth for the first time (for more information on 'selected women', see page 4).

Key findings

- In 2017, 37,913 or 43.1% of selected women giving birth for the first time had an induced labour, an increase from 21,710 or 31.0% in 2004.
- The proportion of Indigenous women who had an induced labour (44.1%) was similar to the proportion of non-Indigenous women who had an induced labour (43.1%).
- The proportion of induced labours in public hospitals was slightly higher compared to private hospitals.


Proportion of selected women giving birth for the first time who had an induced labour, 2004–2007



Notes

1. Data for selected women criteria, parity, were not available from Victoria for 2009.
2. Data for Tasmania from 2004 to 2012 were not available.

Proportion of selected women giving birth for the first time who had an induced labour by hospital sector and Indigenous status, 2017



Women having their second birth vaginally whose first birth was by caesarean section

Caesarean section in a first pregnancy makes a subsequent caesarean section more likely, with 86% of pregnant women with a history of caesarean section birth having a subsequent caesarean section (AIHW 2019).


Both repeat caesarean section (RCS) and vaginal birth after caesarean section (VBAC) have associated risks and benefits. Risks of planned VBAC when compared with planned RCS include haemorrhage, need for blood transfusion, uterine rupture, and perinatal death (Guise et al. 2010, NICE 2019).

This indicator examines women having their second birth vaginally whose first birth was by caesarean section.

Key findings

- In 2017, 3,961 or 12.1% of women who gave birth who had their first birth by caesarean section had a VBAC.
- Rates of VBAC have remained steady from 2007 to 2017 (13.1% and 12.1%, respectively).
- Rates of VBAC were higher in public hospitals and, in 2017, were nearly three times that of private hospitals (15.5% and 5.6%, respectively).
- This is consistent with the overall trend in Australia of private hospitals having a higher rate of caesarean section than public hospitals.
- VBAC was more common among women living in very remote areas (14.5%) than in major cities (11.8%).


Proportion of women whose first birth was by casearean section who had their second birth vaginally, 2007–2017



Notes:

1. Data for Victoria in 2007 to 2009 were not available.
2. Data for Western Australia not published for 2013 as only half a year of data was reported for that year.

Proportion of women whose first birth was by casearean section who had their second birth vaginally, by hospital sector and remoteness of area of usual residence, 2017



General anaesthetic for women giving birth by caesarean section


Regional anaesthesia (or epidural) is the most common method of providing anaesthesia for caesarean section (94%) and is generally safer for the mother and baby than general anaesthesia (AIHW 2019, ANZCA 2018). When general anaesthesia is used, it is typically because of indications such as urgency of the operation, maternal refusal of regional anaesthetic techniques and inadequate or failed regional attempts (Shroff 2004).

This indicator looks at the number of women who received a general anaesthesia when giving birth by caesarean section.

Key findings

- In 2017, 6,367 or 6.1% of women giving birth by caesarean section received general anaesthesia, a slight decrease from 8.2% in 2007.
- Indigenous women giving birth by caesarean section received general anaesthesia more frequently than non-Indigenous women (10.9% and 7.7%, respectively).
- Women living in the most socioeconomically disadvantaged areas were twice as likely to be administered general anaesthetic (8.8%) than women living in the least socioeconomically disadvantaged areas (4.0%).
- Younger women were more likely to receive general anaesthesia for a caesarean section.


Proportion of women having a general anaesthetic when giving birth by caesarean section, by Indigenous status, 2007–2017



Notes

- Caution should be used when interpreting these results. There was no standardised collection across jurisdictions for the data used to derive this indicator.
- Data from Victoria were not available for 2009 or 2010.
- Percentages for Indigenous women are age-standardised.

Proportion of women having a general anaesthetic when giving birth by caesarean section, by mother's age at birth and socioeconomic area of usual residence, 2017



Episiotomy


An episiotomy is an incision made in the perineum (the tissue between the vaginal opening and the anus) and vagina to enlarge the vaginal opening (RCOG 2011). Episiotomy can shorten labour, but the consequences of the trauma to the perineum and, potentially the anus and rectum, need to be balanced against the advantages of an episiotomy. Evidence suggests that selective use of episiotomy is better practice than routine use of episiotomy and may reduce the level of pain, urinary incontinence, painful sexual intercourse or severe perineal trauma (Jiang et al. 2017; RCOG 2011).

This indicator examines the frequency of episiotomy for women having their first baby and giving birth vaginally, either with or without the assistance of instruments.

Key findings

- In 2017, about 1 in 5 women (11,491 or 22.0%) having their first baby received an episiotomy in an unassisted (non-instrumental) birth, an increase from 8,409 or 16.2% in 2004.
- 3 in 4 women (75.2%) received an episiotomy in an assisted (instrumental) birth in 2017, an increase from 60.7% in 2004.
- Episiotomies were slightly more common for women who live in major cities, compared to other regions; this was the case for both assisted and unassisted births.

Proportion of women having their first baby and giving birth vaginally who had an episiotomy, by type of birth (instrumental or non-instrumental), 2004–2017



Notes

1. Caution should be used when interpreting these results. There was no standardised collection across jurisdictions for the data used to derive this indicator.
2. Data from Victoria were not available for 2009.

Birth outcome indicators

Third and fourth degree tears


A perineal tear is the laceration of the skin and tissues that separates the vagina from the anus. Third and fourth degree tears are classified as severe trauma to the perineum involving injury to the anal sphincter muscles and the anal canal. A third or fourth degree tear can occur spontaneously or as a result of obstetric intervention during vaginal birth.

This indicator looks at the proportion of third and fourth degree tears among all women who gave birth vaginally, and among those giving birth vaginally for the first time.

Key findings

- In 2017, third and fourth degree tears were more common in first vaginal births than second or subsequent vaginal births.
- Third and fourth degree tears occurred in 5.0% of first vaginal births and 2.9% of all vaginal births, in 2017.
- Since 2013, the rate of third and fourth degree tears has remained fairly steady for all vaginal births and for all first vaginal births.
- Third and fourth degree tears were more frequent in public hospitals, compared to private hospitals.
- Women having their first vaginal birth who were aged 25 years or older were more likely to experience a third or fourth degree tear than were women in younger age groups.

Proportion of women who had third or fourth degree tears, by mother's age at birth, hospital sector and whether an episiotomy was performed, 2017



Note: For multiple births, the perineal status after the birth of the first-born baby was used.

Apgar score of less than 7 at 5 minutes for births at or after term

Apgar scores are clinical indicators of a baby's condition shortly after birth. The score is based on 5 characteristics of the baby: skin colour, pulse, breathing, muscle tone and reflex irritability and is measured at one and five-minutes after birth. An Apgar score of less than 7 indicates potential complications for the baby and the need for prompt assessment and medical attention (AIHW 2019).


Low Apgar scores can be due to a range of factors, including congenital anomaly, prematurity, perinatal infection, effects of drugs given to the mother for pain relief or anaesthesia, ineffective resuscitation or prolonged hypoxia before birth (Li et al. 2013).

This indicator examines those babies born with an Apgar score of less than 7 at 5 minutes for births at or near term (37 weeks gestation or more).


Key findings

- In 2017, 3,663 or 1.3% of babies born at or near term (37 weeks gestation or more) had an Apgar score of less than 7 at 5 minutes, an increase from 2,229 or 0.9% in 2004.
- Babies born at or near term to women who had labour are slightly more likely to have an Apgar score less than 7 than those born to women who had no labour.
- Babies born to mothers aged younger than 20 were slightly more likely to have an Apgar score less than 7 at 5 minutes than were those born to mothers in older age groups.

Proportion of babies with an Apgar score of less than 7 at 5 minutes for births at or near term (37 weeks gestation), 2004–2017



Proportion of babies with an Apgar score of less than 7 at 5 minutes for births at or near term (37 weeks gestation) by mother's age at birth, 2017



Note: Some caution is required with interpretation of these percentages as the number of babies born at or after term with an Apgar score less than 7 is small.

Small babies among births at or after 40 weeks gestation


Birthweight is a key indicator of infant health, is used as measure of health and wellbeing of the mother in pregnancy, and is an indication of a baby's chance of survival, health, development and wellbeing (AIHW 2016).

This indicator looks at small babies born at or after 40 weeks gestation with a birthweight less than 2,750 grams.

Key findings

- In 2017, 1.3% of babies born after 40 weeks gestation were classified as small.
- The proportion of babies born after 40 weeks gestation who are classified as small has decreased slightly since 2004. The decrease has been most noticeable in babies born to Indigenous women, from 6.1% of babies born after 40 weeks gestation in 2004 to 2.5% in 2017.
- The frequency of small babies born on or after 40 weeks gestation was higher for mothers living in the most disadvantaged socioeconomic areas.
- Babies born at or after 40 weeks gestation were three times as likely to be classified as small if their mother smoked during pregnancy (3.0%) than if their mother did not smoke during pregnancy (1.1%).

Proportion of small babies born at or after 40 weeks gestation by mother's indigenous status, 2004–2017



Proportion of small babies born at or after 40 weeks gestation by mother's socioeconomic area, 2017



Notes

- Caution should be used when comparing these results with other nationally and internationally reported indicators, as the cut-off birthweight may be different to this indicator.
- Some caution is required with interpretation of these percentages as the number of small babies born at or after 40 weeks gestation is small.
- Percentages for Indigenous women are age-standardised.

Data source and methods

The data used for the National Core Maternity Indicators is sourced from the National Perinatal Data Collection (NPDC). The NPDC is a national population-based cross-sectional collection of data on pregnancy and childbirth. The data are based on births reported to the perinatal data collection in each state and territory in Australia. Midwives and other birth attendants, using information obtained from mothers and from hospital or other records, complete notification forms for each birth.

The NPDC consists of the Perinatal National Minimum Data Set (NMDS) and some additional data items. With the exception of two indicators (Apgar score and small babies at term), the indicator definitions rely to some extent on non-NMDS data items, which are provided by jurisdictions on a voluntary basis and may be affected by differences in collection practices.

Information is not available from all jurisdictions to support the 12 indicators in all years. An overview of data availability for reporting against relevant indicators can be found at <<https://www.aihw.gov.au/reports/mothers-babies/ncmi-data-visualisations/contents/technical-notes/data-source-and-quality>>.

More information

For more information on these indicators, see the online report at:
<<https://www.aihw.gov.au/reports/mothers-babies/ncmi-data-visualisations/contents/summary>>.

Acknowledgments

This report was written by Madeline Elkington and Jacqueline Rek of the Australian Institute of Health and Welfare. The authors gratefully acknowledge the contributions of the following AIHW staff: Tracie Reinten, Joshua Alexander, Amanda Reyes-Veliz, Bernice Cropper, Fadwa Al-Yaman, Michael Frost, Michael Humphrey and Elizabeth Ingram.

State and territory health departments are thanked for their contribution to the NPDC and their review of this report.

References

- ACSQHC (Australian Commission on Safety and Quality in Health Care and Australian Institute of Health and Welfare) 2018. The Third Australian Atlas of Healthcare Variation. Sydney: 2018
- AIHW (Australian Institute of Health and Welfare) 2016. Australia's health 2016. Australia's health series no. 15. Cat. no. AUS 199. Canberra: AIHW.
- AIHW 2018. Australia's mothers and babies 2016—in brief. Perinatal statistics series no. 34 Cat. No. PER 97. Canberra: AIHW.
- AIHW 2019. Australia's mothers and babies 2017—in brief. Perinatal statistics series no. 35 Cat. No. PER 100. Canberra: AIHW.
- ANZCA (Australian and New Zealand College of Anaesthetists) 2018. Risks and complications of anaesthesia. Viewed August 2019, <<http://www.anzca.edu.au/patients/frequently-asked-questions/risks-and-complications>>.
- Betran A, Ye J, Moller A, Zhang J, Gürmezoglu A & Torloni M 2016. The increasing trend in caesarean section rates: global, regional and national estimates: 1990–2014. Public Library of Science 11(2):e0148343.
- DoH (Department of Health) 2018. Clinical Practice Guidelines: Pregnancy Care. Canberra: Australian Government Department of Health.
- Guise JM, Eden K, Emeis C, Denman MA, Marshall N, Fu RR et al. 2010. Vaginal birth after caesarean: new insights. Evidence Report Technology Assessment (Full Report) 191:1–397.
- Jiang H, Qian X, Carroli G & Garner P 2017. Selected versus routine use of episiotomy for vaginal birth. Cochrane Database of Systematic Reviews 2: CD000081.
- Keag O, Norman J & Stock S 2018. Long-term risks and benefits associated with cesarean delivery for mother, baby, and subsequent pregnancies: Systematic review and meta-analysis. Public Library of Science: Medicine 15(1):e1002494.
- Li F, Wu T, Lei X, Zhang H, Mao M & Zhang J 2013. The Apgar score and infant mortality. Public Library of Science 8(7):e69072.
- McDonnell R 2011. Induction of labour. O&G Magazine 13(3):62–4.
- NICE (National Institute for Health and Care Excellence) clinical guideline CG132; Caesarean section: 2019.
- RANZCOG (The Royal Australian and New Zealand College of Obstetricians and Gynaecologists) 2016. Instrumental vaginal birth: Position statement (PDF). Viewed August 2019, <[https://ranz cog.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Instrumental-Vaginal-Birth-\(C-Obs-16\)-Review-March-2016.pdf?ext=.pdf](https://ranz cog.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Instrumental-Vaginal-Birth-(C-Obs-16)-Review-March-2016.pdf?ext=.pdf)>

RCOG (The Royal College of Obstetricians and Gynaecologists) 2011. Operative vaginal delivery: Green top guideline no. 26 (PDF). Viewed August 2019, <https://www.rcog.org.uk/globalassets/documents/guidelines/gtg_26.pdf>.

Rowland I & Redshaw M 2012. Mode of birth and women's psychological and physical wellbeing in the postnatal period. *BMC Pregnancy and Childbirth* 12:138.

Shroff R, Thompson A, McCrum A & Rees S 2004. Prospective multidisciplinary audit of obstetric general anaesthesia in a district general hospital. *Journal of Obstetrics and Gynaecology* 6:641–646.

Villar J, Carroli G, Zavalenta N, Donner A, Wojdyla D, Faundes A et al. 2007. Maternal and neonatal individual risks and benefits associated with caesarean delivery: multicentre prospective study. *British Medical Journal* 335:1025.

WHO (World Health Organization) 2013. WHO recommendations for the prevention and management of tobacco use and second-hand smoke exposure in pregnancy. Viewed August 2019, <https://apps.who.int/iris/bitstream/handle/10665/94555/9789241506076_eng.pdf;jsessionid=F48A29577A24E7FE2478879F1524D783?sequence=1>.



The National Core Maternity Indicators (NCMIs) present information on measures of clinical activity and outcomes to assist in improving the quality of maternity services in Australia by establishing baseline data for monitoring and evaluating practice change. These indicators cover data for the majority of women who gave birth in Australia from 2004 to 2017 and are grouped into 3 broad topic areas—antenatal period, labour and birth and birth outcomes.

aihw.gov.au



Stronger evidence,
better decisions,
improved health and welfare